

University of Groningen

The role of vitamin A in bile acid synthesis and transport and the relevance for cholestatic liver disease

Hoeke, Martijn Oscar

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:
2013

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Hoeke, M. O. (2013). *The role of vitamin A in bile acid synthesis and transport and the relevance for cholestatic liver disease*. s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Chapter 1

The Interrelationship Between Bile Acid and Vitamin A Homeostasis

Martijn O. Hoeke and Klaas Nico Faber

Department of Gastroenterology and Hepatology, University Medical Center Groningen,
Groningen, University of Groningen, The Netherlands

ABSTRACT

This thesis explores the regulatory interaction between vitamin A and bile salts. Bile salt homeostasis is of crucial importance to maintain vitamin A homeostasis. Bile salts are the emulsifying components of bile that are required for the effective absorption of the fat-soluble vitamin A in the intestine, most of which is stored in stellate cells in the liver. Dysregulation of bile salt homeostasis, for instance in patients with cholestasis, causes vitamin A malabsorption. At the same time cholestatic liver disease promotes the release of vitamin A from hepatic stellate cells, promoting depletion of vitamin A from the body. A functional relationship in the opposite direction, in which vitamin A influences bile salt homeostasis, has only recently emerged. Complex regulatory pathways ensure that bile salt homeostasis is maintained. These processes involve bile acid sensors, which are ligand-activated transcription factors. This way, bile salts regulate their own synthesis and transport at the transcriptional level. These bile acid sensors need to physically interact with a vitamin A sensor in order to regulate transcription. In this thesis, we show that vitamin A is required to maintain bile salt homeostasis and protects the liver during obstructive cholestasis. This chapter introduces the reader to the origin and function of bile salts and vitamin A and the signalling pathways they control.

1. THE ENTEROHEPATIC CIRCULATION OF BILE ACIDS

The liver is the largest solid organ of the human body and well-known for its regenerative capacity. The liver consists of various cell types, including hepatocytes, stellate cells, cholangiocytes, endothelial cells, Kupffer cells (hepatic macrophages), and oval or progenitor cells, each with their specific role in the function of the liver. Hepatocytes (the hepatic parenchymal cells) make up 80% of the liver mass and fulfil most of the primary functions of the liver. Hepatocytes are arranged in plates that form a physical barrier between the blood in the sinusoid and the bile in the canaliculus. Approximately 80% of the hepatic blood supply enters via the portal vein and is rich in nutrients originating from the intestine. Oxygen-rich blood enters via the hepatic artery, mixes with the portal blood and flows along the hepatic sinusoids and exits via the central vein. The sinusoids (**Figure 1.1**) are lined with a sheet of endothelial cells, that allows for the exchange of small molecules between blood and hepatocytes. Kupffer cells are integrated into the sheet of endothelial cells. Stellate cells reside in the space of Disse, the plasma-filled space between hepatocyte and sinusoid. Stellate cells (Ito cells, fat-storing or vitamin A-storing cells) are myofibroblast-like cells that contain large cytoplasmic lipid droplets. Cholangiocytes are the bile duct epithelial cells and form approximately 3% of the liver mass (1).

In broad terms, the physiological functions of the liver can be described as, synthesis, storage, metabolism, catabolism and excretion. Proteins synthesized by the liver include serum proteins such as albumin, different coagulation factors (such as fibrinogen) and the retinol binding protein 4 (RBP4).

Cholesterol and triglycerides are synthesized by the liver. Moreover, it is the main storage site of glucose (in the form of glycogen), iron and vitamin A. Another important function of the liver is detoxification. Endogenous waste products (such as bilirubin) and exogenous harmful compounds (drugs and toxins) are metabolized by the liver. This makes them less toxic and increases water solubility, which enhances clearance of these compounds. The liver is also the largest gland of the human body, amongst others synthesizing and excreting the hormone insulin-like growth factor 1 (IGF-1) and producing bile. Bile aids in the digestion and absorption of nutrients and is the main route for the excretion of waste products via the intestine. Major components of bile are bile salts, phospholipids, cholesterol, bilirubin, alkaline hydrogen carbonate ions (HCO_3^-) and water. Bile salts are the components of the bile that give it its fat-emulsifying properties. Bile salts are synthesized by hepatocytes using cholesterol as substrate. The liver secretes bile via the bile ducts and the gallbladder into the intestinal lumen (duodenum). Here, bile fulfils its function in the digestion and absorption of fats and fat-soluble dietary compounds. In the terminal ileum, bile salts are reabsorbed and transported back to the liver via the portal vein. This shuttling of bile salts between liver and intestine is called the enterohepatic circulation of bile salts and is a very efficient process as per cycle only 5% of the bile salts are lost in the feces. *De novo* synthesis of bile salts in the liver compensates for this loss and maintains a balanced amount of bile salts in the enterohepatic cycle.




Figure 1.1. Structure of the liver sinusoid

The liver possesses a dual blood supply. Nutrient-rich blood coming from the portal vein and oxygen-rich blood coming from the hepatic artery merge upon entering the sinusoid. The sinusoids are lined with endothelial cells and Kupffer cells. The plasma-filled space of Disse separates the endothelial cells from the hepatocytes. Bile produced by the hepatocytes is excreted into the bile canaliculi that empty into the bile ductule, which is formed by cholangiocytes. Vitamin A-storing stellate cells reside within the space of Disse. Reprinted from Frevert (2004) (361).

1.1. Bile Acid/Salt Biosynthesis

Bile acid synthesis produces two main types of primary bile acids: cholic acid (CA) and chenodeoxycholic acid (CDCA). These are conjugated to either glycine or taurine, yielding the bile salts taurocholic acid (TCA), glycocholic acid (GCA), taurochenodeoxycholic acid (TCDCA) and glycochenodeoxycholic acid (GCDCA). Bile salts that are not absorbed at the terminal ileum move to the colon where resident bacteria may convert them to secondary bile acids, such as lithocholic acid (LCA) and deoxycholic acid (DCA). A significant fraction of these secondary bile acid are also absorbed into the circulation. DCA accumulates in the bile acid pool, LCA to a much lesser extend as it is sulfated and subsequently excreted into bile (reviewed in (3)).

The term bile “acid” refers to the protonated ($-\text{COOH}$) form, while bile “salt” refers to the deprotonated or ionized ($-\text{COO}^-$) form. Bile acids conjugated to glycine or taurine are largely deprotonated at the physiological pH, and should thus be regarded as bile salts. The ionized conjugated bile salt is more hydrophilic when compared to the unconjugated bile acid. Conjugated bile acids are therefore more efficient in emulsifying fats than their unconjugated counterparts.

Cholesterol is oxidized to bile acids through a cascade of enzymatic conversions involving at least 13 different enzymes (see for a review (4)). Although the first steps of the conversion from cholesterol to bile acids may take place at extra-hepatic locations, the production of the end products, e.g. bile salts, is restricted to the hepatocytes in the liver. Two enzymes, cholesterol 7- α -hydroxylase (CYP7A1) and sterol 12- α -hydroxylase (CYP8B1) are of particular relevance. CYP7A1 is considered the rate-limiting enzyme in bile acid synthesis (5), while CYP8B1 converts CDCA to CA and thus determines the relative hydrophobicity of the bile acid pool (6). The final step in bile acid synthesis is the conjugation of taurine or glycine and is exclusively performed by the peroxisomal enzyme bile acid coenzyme A: amino acid N-acyl-transferase (BAAT) (7). Bile acid synthesis is increased following feeding and glucose and insulin are important factors that induce CYP7A1 gene expression (8, 9).

1.2. Transporter Proteins Involved in (Bile Formation and) Bile Salt Transport

Once conjugated, membrane translocation of bile salts requires active transport by membrane-imbedded transporters. These transporters reside in the membranes of polarized hepatocytes, cholangiocytes and enterocytes and keep the bile salts in the enterohepatic circulation. The transporter proteins involved in the enterohepatic circulation have been the subject of multiple reviews (10-13). A schematic overview of the enterohepatic circulation of bile salts and the transporter proteins involved is depicted in **Figure 1.2**.

1.2.A. Hepatic Bile Salt Transport Mechanisms

Hepatic Basolateral Transport/Bile Salt Import

Bile salts returning from the intestine via the blood circulation are absorbed into the hepatocyte by the sodium/taurocholate co-transporting polypeptide (NTCP) (15, 16)

and at least three members of the organic anion transporting polypeptide (OATP) family (17). A recent study showed that also unconjugated bile acids are subject to active transport via OATP1B2 rather than entering the cell by passive diffusion (18). NTCP (SLC10A1) belongs to the SLC10 family, which comprises transport proteins that transport organic salts in a sodium-dependent manner. The human NTCP protein consists of 349 amino acids with a molecular mass of 56 kDa. In the liver, NTCP is expressed exclusively at the basolateral membrane of the hepatocyte and preferentially transports taurine conjugated bile acids. (15, 19, 20). However, also various other physiological bile salts and the steroid conjugate estrone 3-sulfate are substrates of NTCP (21). Expression of NTCP in pancreatic acinar cells has also been reported (22).

The superfamily of OATPs comprises a family of transporters with a broad spectrum of substrates. The substrate specificity of OATPs is reviewed in detail by König *et al.* (23). While most of the OATP proteins are expressed in multiple tissues, OATP1B1 (OATP2/OATP-C/SLCO1B1/SLC21A6) and OATP1B3 (OATP8/SLCO1B3/SLC21A8) are predominantly if not exclusively expressed in liver at the basolateral membrane of hepatocytes. OATP1B1 consists of 691 amino acids with a molecular mass of 84 kDa. OATP1B3 consists of 702 amino acids and represents a glycoprotein with a molecular mass of 120 kDa. Both of these OATPs were shown to transport tauro- and glyco-conjugated CA (10, 23-25). OATP1A2 (OATP-A/SLCO1A2) is mainly expressed in brain and liver. It has a molecular mass of 85 kDa and consists of 670 amino acids. The rodent ortholog is OATP1A1, which has 12 putative transmembrane spanning domains. OATP1A2 is localized to the basolateral membrane of the hepatocyte and is reported to transport both conjugated and unconjugated CDCA and CA (10, 26, 27).

Hepatic Intracellular Transport of Bile Acids

The cytosolic liver fatty-acid-binding protein (L-FABP) is involved in fatty acid metabolism and transport. Moreover, L-FABP was found to bind bile acids in a 1:2 ratio and to be excreted into bile (28, 29). Although bile acid metabolism is altered in *L-fabp*^{-/-} mice, the physiological role of L-FABP in bile acid trafficking remains to be determined (30).

Part of the bile salt pool that enters the liver is deconjugated by intestinal flora when passing through the intestine. Before excretion into bile, *de novo* and recycled bile acids need to be (re-)conjugated. The enzyme responsible for bile acid conjugation (BAAT) resides solely in peroxisomes, which suggests the existence of intracellular bile acid/salt transporters. Although no intracellular transporter has been identified to date, recent data from our laboratory strongly supports the existence of such bile salt transporters in the peroxisomal membrane (4, 7, 31).




Figure 1.2. Transport mechanisms involved in the enterohepatic circulation of bile salts and formation of bile

The enterohepatic circulation of bile salts (BSs) requires dedicated transporter proteins on both sides of polarized hepatocytes, cholangiocytes and enterocytes. Bile salts returning from the intestine via the blood circulation are absorbed into the hepatocyte by the sodium/taurocholate co-transporting polypeptide (NTCP) and at least three members of the organic anion transporting polypeptide (OATP) family. Conjugated monovalent BSs are excreted from the hepatocyte via the canalicular bile salt export pump (BSEP). Bivalent BSs, together with organic anions (OA⁻), are excreted via the canalicular multidrug resistance-associated protein 2 (MRP2). The ABC half-transporters ABCG5/G8 secrete cholesterol into bile. The multidrug resistance protein 3 (MDR3) and the familial intrahepatic cholestasis type 1 (FIC1) protein excrete phosphatidylcholine (PC) and phosphatidylserine (PS), respectively, which form mixed micelles in bile together with cholesterol and BS⁻. Organic cations (OC⁺) are excreted by the efflux pump for xenobiotic compounds (MDR1). Cholangiocytes and enterocytes use similar transport mechanisms for BS⁻ shuttling. At the apical membrane, BS⁻ are taken up by the ileal bile salt transporter (ASBT). Cellular transport of BS⁻ in the enterocyte is facilitated by the ileal bile acid binding protein (IBABP). Excretion of BS⁻ is primarily performed by the organic solute transporter dimer (OSTα/β). Besides excreting OA⁻, the multidrug resistance-associated protein 3 (MRP3) provides an alternative route for the elimination of BS⁻. Both OSTα/β and MRP3 are also expressed at

the basolateral membrane of hepatocytes, in particular in pathological conditions. Here they provide an alternative route for the elimination of BS⁻ and other OA⁻. MDR1 is also present in the cholangiocytes and enterocytes at the apical membrane, while MRP2 is also expressed in ileum. BS⁻ excreted by the enterocytes/cholangiocytes return to the liver via the blood, closing the enterohepatic circulation. Inspired by Trauner *et al.* (2003) (11) and Zollner *et al.* (2006) (276).

Hepatic Canalicular Bile Salt Export

Conjugated recycled and *de novo* synthesized bile salts are excreted into bile by the bile salt export pump (BSEP/ABCB11). The 160 kDa BSEP-protein is exclusively expressed at the canilicular membrane of hepatocytes and belongs to the B subfamily of ATP-binding cassette (ABC) transporters. A total of 48 ABC type transporters have been identified in the human genome. BSEP contains 12 membrane spanning domains and two intracellular loops that contain nucleotide/ATP binding motifs, which is the typical build-up of an ABC transporter. Mutations in the *BSEP* gene are associated with progressive familial intrahepatic cholestasis type 2 (PFIC2), which is characterized by severe and progressive liver disease. In addition, mutations in *BSEP* may cause benign recurrent intrahepatic cholestasis (BRIC) that is characterized by intermittent attacks of cholestasis without permanent liver damage (32-37).

The multidrug resistance-associated protein 2 (MRP2/ABCC2) transports a variety of organic anions at the canalicular membrane into bile, including conjugated bilirubin (38). MRP2 belongs to the C subfamily of ABC transporters (MRPs), of which ten members (*MRP1* to 9 and pseudo gene *ABCC13*) have been identified (39). Besides the liver, MRP2 is expressed in the intestine (duodenum) and kidney (40). MRP2 is capable of transporting bivalent glucuronidated and sulfated bile salts, and was recently also shown to transport tauroursodeoxycholate (41). The role of MRP2 in maintaining bile salt homeostasis may differ between species. Mutations in *MRP2* lead to Dubin-Johnson syndrome in humans (42), a benign form of jaundice (43, 44), transport of bile salts into bile is not disturbed in these patients. However, Groningen yellow (GR/TR) rats, which lack a functional MRP2 protein, display an impaired bile flow (45).

Additional Hepatic Canalicular Transporter Proteins That Contribute to Bile Formation

The multidrug resistance protein 3 (MDR3/ABCB4; the rodent ortholog is MDR2) and the ABCG5/ABCG8 dimer transport phospholipids and cholesterol, respectively, from the inner to the outer leaflet of the canalicular membrane. Bile salts excreted into the canaliculus interact with the outer leaflet of the hepatocyte's apical membrane and extract cholesterol and phospholipids thereby forming mixed micelles (46). Like BSEP, MDR3 belongs to the B subfamily of the ABC transporter family and shares the 12 membrane spanning domain structure. Defects in the *MDR3* gene cause progressive familial intrahepatic cholestasis type 3 (PFIC3). In this disease, phospholipid transfer to bile is disturbed. Lack of phospholipids hampers the formation of the mixed micelles, increasing the concentration of pure micelles in bile, resulting in damage to the bile duct epithelium and cholestasis (47, 48).

Another member of the B subfamily of ABC transporters is *ATP8B1*, it encodes FIC1, named after the associated disease familial intrahepatic cholestasis type 1 that is caused by mutation in the *ATP8B1* gene (reviewed in (49)). The FIC1 protein acts as a flippase for phosphatidylserine. Although FIC1 does not transport bile salts, mutations in the *ATP8B1* gene cause cholestasis. The proposed mechanism for this observation is that in the absence of functional FIC1 protein the membrane gets depleted of lipids (including cholesterol) in the process of bile formation. Lack of cholesterol in the membrane limits the function of BSEP resulting in intracellular bile salt accumulation (49).

ABCG5 and ABCG8 are members of the G subfamily of the ABC-transporters. They are so-called half-transporters that form a heterodimer to become a functional cholesterol transporter. ABCG5/G8 is localised to the apical membrane of hepatocytes, and cholangiocytes and enterocyte (50, 51). Defects in either the murine *Abcg5* or *Abcg8* gene are associated with decreased excretion of cholesterol in bile and an increased absorption of cholesterol in the intestine (52).

The multidrug resistance-1 P-glycoprotein (MDR1/ABCB1) is an efflux pump for xenobiotic compounds and has a broad substrate specificity. It is associated with the resistance of tumor cells against chemotherapeutic drugs. MDR1 is expressed at the canalicular membrane of hepatocytes and the apical side of the bile ductal and intestinal epithelium (53). Apart from metabolites of LCA, bile salts do not seem to be substrates for MDR1 and malfunctioning of MDR1 does not result in cholestatic liver disease (54). However, accumulation of certain drugs and substrates for MDR1 may inhibit the function of BSEP (12, 55). Unlike human PFIC2 patients, *Bsep*^{-/-} mice do not suffer from severe cholestasis and have substantial bile salt secretion (56). Interestingly these mice have increased MDR1A expression while expression of MDR2, MRP2, and MRP3 (the latter will be discussed later) was only slightly increased (57). This suggests that MDR1A in mice might be involved in bile salt transport in the absence of BSEP.

1.2.B. Bile Ductular and Intestinal Bile Transport Mechanisms

After hepatobiliary secretion, bile salts travel through the bile ducts and are stored in the gallbladder. Triggered by food ingestion, the gallbladder contracts and bile is released into the duodenum where bile salts fulfil their function in the digestion and absorption of fat-soluble nutrients. At the terminal ileum, they are efficiently reabsorbed to the circulation and returned to the liver via the portal tract. The bile ductular and distal ileum epithelial cells, cholangiocytes and enterocytes, contain highly similar bile salt transport mechanisms (**Figure 1.2**).

Apical Sodium-dependent Bile Salt Import

The apical sodium-dependent bile salt transporting protein (ASBT/SLC10A2) was initially identified as the intestinal analogue of Ntcp, belonging to the same SLC10 family of sodium-dependent organic ion transporters. Subsequently, ASBT was also shown to facilitate bile salt absorption at the apical domain of rat cholangiocytes.

The membrane topology and transport characteristics are similar to NTCP. The ASBT glycoprotein consists of 348 amino acids and has a molecular mass of 48 kDa (12, 19, 58-60). Mutation in the *SLC10A2* gene have been associated with primary bile salt malabsorption and patients suffer from diarrhea and fat malabsorption (61).

Intracellular Bile Salt Transport

The intestinal bile acid binding protein (IBABP/FABP6) is expressed in enterocytes and cholangiocytes. IBABP represents a protein of 128 amino acids with a molecular mass of 14-15 kDa. IBABP is proposed to be the most important protein for transcellular bile salt movement in enterocytes. However, its physiological function remains elusive, as the absence of IBABP, as observed in *Fxr*^{-/-} mice, does not affect the turnover rate and cycling time of the cholate pool (62). Recent data obtained with *Fabp6*^{-/-} mice show that IBABP is not prerequisite for ileal bile acid absorption but that it does contribute to efficient shuttling of bile acids through the enterocyte (63). Additionally, binding of bile salts by IBABP has also been suggested to prevent intracellular toxicity (12, 64, 65).

Basolateral Bile Salt Efflux Mechanisms

The identity of the transporter that facilitates the efflux of bile salts from ileal enterocytes across the basolateral membrane to the circulation has long been unclear. Initially, it was proposed that this activity was performed by a truncated form of ASBT (t-ASBT) that is a product of alternatively splicing of the corresponding transcript. T-ASBT was shown to be capable of transporting TCA *in vitro* and to be expressed at the basolateral membrane. T-ASBT was identified in ileum, kidney, and bile ductal epithelial cells but not in hepatocytes (12, 66). A second transporter that was proposed to export bile salts from enterocytes was the multidrug resistance-associated protein 3 (MRP3/ABCC3). MRP3 is expressed in various tissues of the gastrointestinal tract, including liver and ileum, where it was shown to be expressed at the basolateral membrane of cholangiocytes and enterocytes, respectively. MRP3 was shown to be capable of transporting various endogenous bile salts. However, mice lacking *Mrp3* do not show altered bile salt physiology. This is in line with the observation that MRP3 expression is rather low in human liver under healthy conditions (39, 67-71). With the identification in 2005 of the organic solute transporter alpha/beta (OST α/β) as the main functional basolateral bile salt transporter in the mouse ileum, the missing link in bile salt circulation was identified (72, 73). OST α/β consists of an α - and a β -subunit that together form a functional transporter at the basolateral membrane. OST α/β was initially identified as an organic solute and steroid transporter in the marine vertebrate little skate (74). OST α is a 340 amino acid protein, and is predicted to contain seven transmembrane domains. The β -subunit is a much smaller protein of 128 amino acids predicted to have a single transmembrane domain (75). The human/rodent homologs were found to export various endogenous bile salts. OST α/β is predominantly expressed at the basolateral membrane of intestinal epithelium and cholangiocytes in liver, with less but detectable amounts also in renal proximal tubu-

lar cells in kidney and hepatocytes in the liver (76). Interaction between the α - and β -subunits is essential for protein stability and efficient trafficking through the Golgi apparatus to the plasma membrane. *Osta*^{-/-} mice have normal *Ost* β mRNA levels, but the OST β protein is absent (77). Emerging evidence suggests a pleiotropic function of OST α/β that is not limited to bile salt transport (78). Bile salts that are excreted to the circulation by the enterocytes are transported back to the liver, closing the circle of the enterohepatic circulation.

1.3. The Adaptive Response to Bile Salt Accumulation

Bile salt homeostasis is maintained by coordinated regulation of bile salt synthesis and transport. When bile acid concentrations increase, for instance when mice are fed a bile salt-containing diet, expression of both *Cyp7a1* and *Cyp8b1* is reduced, which represses bile acid synthesis. At the same time expression of *Ntcp* is lowered and *Bsep* expression is increased (79). In obstructive cholestasis, as in patients with primary biliary cirrhosis (PBC) and in bile duct-ligated rodents, hepatic expression of OST α , OST β , MRP3 and multidrug resistance-associated protein 4 (MRP4/ABCC4) is increased (80-83). Intestinal expression of *ASBT* mRNA is lowered by elevated bile salt levels (84). The fact that *Mrp4*^{-/-} mice have an impaired cytoprotective response in obstructive cholestasis exemplifies that these adaptive responses are crucial to prevent excessive liver damage when bile salt flow from the liver is impaired (83). Although MRP3 was shown to transport (secondary) bile salts *in vitro*, bile duct-ligated *Mrp3*^{-/-} mice do not suffer more liver damage than wild type mice (69). In this case MRP4 may be sufficient to prevent excessive liver damage. These mechanisms protect the hepatocyte and enterocyte from cytotoxic accumulation of bile acids. Regulation of these adaptive mechanisms will be discussed later.

2. VITAMINS, ESSENTIAL MICRONUTRIENTS

In 1916, vitamin A was one of the first vitamins that was discovered (85). Historically, the word “vitamine” was used to describe a growth factor that is present in food and that is essential for life. The combination of the words “vital” and “amine” yielded “vitamine”. As it became apparent that there were multiple vitamins, McCollum divided them into two classes: “fat-soluble factor A” and “water-soluble factor B”. Following the discovery that not all of these micronutrients shared the amine-group in their molecular structure, Drummond changed the name to “vitamin”, dropping the final “e”. He renamed McCollum’s compounds into vitamin A and vitamin B and new members of these series of factors were named vitamin C, vitamin D etc. (85, 86). Water-soluble vitamins are readily expelled from the body, hence, regular (daily) intake is required. In contrast, fat-soluble vitamins are stored in the liver and adipose tissues. Stable levels of these vitamins can therefore be maintained for a longer time period. The downside of this is that excessive intake of the fat-soluble vitamins A, D, E and K poses a greater risk of vitamin intoxication than vitamins B and C. The uptake of fat-soluble vitamins requires the presence of bile salts in the intestinal lumen. Cholestatic liver diseases, such as PBC, primary sclerosing cholangitis (PSC), biliary

atresia and gallstone disease, which dysregulate the flow of bile to the intestine, may lead to malabsorption of fat-soluble vitamins and associated clinical symptoms of hypovitaminosis.

The primary focus of this thesis is the fat-soluble vitamin A, but also the other fat-soluble vitamins will be briefly discussed.

2.1. Vitamin A

In 1930, Moore proved that the yellow plant pigment beta-carotene is converted to colourless retinol in the mammalian body (87). In 1931, Karrer deduced the molecular structure of retinol and beta-carotene (85, 86) (**Figure 1.3A-B**). Nowadays, the term vitamin A is used as the generic descriptor for retinoids exhibiting the biological activity of retinol. Retinoids are a class of compounds consisting of four isoprene (2-methyl-1,3-butadiene) units joined in a head-to-tail manner. (86).

Vitamin A-derivatives fulfil numerous important functions in the mammalian body, including roles in vision, maintenance of epithelial surfaces, immune competence, reproduction and embryonic growth and development (88). Dietary sources of vitamin A are provitamin A carotenoids (mainly β -carotene, from plant sources), preformed vitamin A (retinyl esters from animal sources) and precursors of retinol (88). The recommended daily intake of vitamin A is approximately 700 and 900 μg for adult women and men, respectively. Dietary intake of solely β -carotene may be inadequate to maintain normal levels of vitamin A, retinyl esters should therefore be considered an essential component of a healthy diet (89). Approximately 80% of the total body pool of vitamin A is stored in the liver as retinyl esters (90).

Some of vitamin A's numerous biological functions are related to its antioxidant properties. Especially carotenoids, such as lycopene and β -carotene, are potent antioxidants (91, 92), but also retinol, retinyl esters and retinoic acid (RA) have been shown to prevent lipid peroxidation *in vitro* (93, 94). Another well-known function of vitamin A is its role in the visual cycle by photoisomerization. Rhodopsin, the visual pigment of the rod photoreceptor cell, contains 11-cis retinal as its light-sensitive cofactor. Light activation is achieved by 11-cis to all-trans isomerisation, followed by the release of all-trans retinaldehyde (95).

However, most of the biological functions of vitamin A involve the activation of ligand-dependent transcription factors. This hormonal function of vitamin A gained tremendous scientific interest with the discovery of two vitamin A receptors that are members of the nuclear receptor superfamily (96). All-trans retinoic acid (atRA) and 9-cis retinoic acid (9cRA) (**Figure 1.3C-D**) are the two biologically active isomers of RA that regulate gene transcription via the retinoic acid receptor (RAR) and the retinoid X receptor (RXR). These nuclear receptors will be discussed in section 4.




Figure 1.3. Fat soluble vitamins A, D, E and K

Beta- (β -) carotene (A) is the primary source of vitamin A in plant-derived foods, which can be cleaved into two molecules of retinol (B). Retinol and its esters are the main transport and storage forms of vitamin A. The 2 predominant biologically active forms of vitamin A are *all-trans* and 9-*cis* retinoic acid (C and D). The biologically active compound of vitamin D is calcitriol (1,25-dihydroxycholecalciferol or 1,25-dihydroxyvitamin D₃) (E). Alpha- (α -) tocopherol (F) is the most abundant form of vitamin E in nature, it also has the highest biological activity. Phyloquinones (G) are the primary dietary source of vitamin K, while menaquinones (H) are produced by bacteria in the gastro-intestinal tract.

2.2. Vitamin D, Important Micronutrient But Not Really a Vitamin

Vitamin D was discovered as an essential nutrient to prevent rickets, e.g. softening of bones leading to skeletal deformations in children and an increased risk of fractures later in life. Vitamin D deficiency can result in osteoporosis (decreased bone mineral density) in adults, causing osteomalacia (softening of bone), muscle weakness and increased risk of fractures (97). Vitamin D is required for optimal absorption of dietary calcium and phosphate and plays an essential role in calcium homeostasis

and bone metabolism. UV-light irradiation of the skin was found to prevent rickets as this stimulates formation of vitamin D₃ in the skin. The ability to produce sufficient amounts of vitamin D₃ with adequate sunlight exposure indicates that vitamin D is actually not a vitamin (98). The name “vitamin D” refers to a group of steroids that exhibit the biological activity of cholecalciferol. The two physiologically relevant forms are vitamin D₂ (ergocalciferol) and vitamin D₃ (cholecalciferol). Very few foods contain significant amounts of vitamin D. D₂ is plant derived and is present in mushrooms that are exposed to UV-light. The animal-derived D₃ is present in, amongst others, fish oil, egg yolk and liver. (97, 99).

In the liver, vitamin D₃ is metabolized to 25-hydroxycholecalciferol (calcidiol, 25-(OH)D), which is subsequently metabolized by the kidney into the biologically active steroid hormone 1,25-dihydroxyvitamin D [1,25(OH)₂D] (**Figure 1.3E**) (97). Vitamin D is stored in adipose tissues as calcidiol. Hypovitaminosis D may develop through a combination of inadequate vitamin D intake and insufficient exposure to sun light in otherwise healthy people (97, 100). In addition, vitamin D deficiency may develop as a result of multiple disease conditions, including impaired synthesis in the skin, impaired intestinal absorption as well as acquired and heritable disorders of vitamin D metabolism and responsiveness.

2.3. Vitamin E, Solely Antioxidant or More

In 1922, vitamin E was first described as an essential nutrient for reproduction in rats. Vitamin E is the generic description for α -, β -, γ -, and δ - tocopherols and the corresponding four tocotrienols. Alpha-tocopherol is the most abundant form in nature, which has the highest biological activity and reverses symptoms of vitamin E deficiency in humans (**Figure 1.3F**) (101-103). Nuts, seeds, and vegetable oils are among the richest sources of alpha-tocopherol and significant amounts are present in green leafy vegetables. Tocopherol is a potent antioxidant that prevents lipid peroxidation of polyunsaturated fatty acids in cellular membranes. In addition to its antioxidant activities, vitamin E is reported to be involved in cell signalling, immunity, gene expression and metabolic processes (104). However, others believe that all vitamin E actions arise from its antioxidant protection of polyunsaturated fatty acids in the plasma membrane, and that vitamin E is nothing more than an antioxidant (105). The liver plays a central role in α -tocopherol homeostasis. Vitamin E travels the body as a component of lipoproteins that are taken up (chylomicrons, low density lipoprotein (LDL) and high density lipoprotein (HDL)) and excreted (very low density lipoprotein (VLDL)) by the liver. One third of the body content of vitamin E is stored in the liver, mainly in hepatocytes (106).

2.4. Vitamin K

Vitamin K refers to a group of chemically similar compounds called naphthoquinones. The naturally occurring forms of vitamin K are K₁- (phyloquinone) and the K₂-compounds (menaquinones) that contain unsaturated side chains of varying length (**Figure 1.3G-H**). Phytonadione is found in plants and is the primary

dietary source of vitamin K for humans. Menaquinones are produced by bacteria in the gastrointestinal tract and contribute to the intake of vitamin K (107-109).

Vitamin K is required for normal blood clotting. In fact, the name vitamin K is derived from “Koagulations-Vitamin” in German and Scandinavian languages (110). It acts as a cofactor for γ -glutamyl carboxylase. This enzyme is responsible for post-translational modification of specific glutamate residues in proteins to γ -carboxyglutamate. The majority of γ -carboxylated proteins function in blood coagulation, while others play a role in calcium homeostasis and normal bone metabolism (111, 112).

Green vegetables, such as Brussels sprouts, spinach, cauliflower and broccoli, as well as certain vegetable oils, including soybean oil, rapeseed oil and olive oil, are rich dietary sources of vitamin K. Animal-derived foods contain limited amounts of vitamin K (109).

Vitamin K deficiency occurs when intake is inadequate or when the intestinal flora is altered. Clinical symptoms of vitamin K deficiency include clotting disorders (e.g. vitamin K deficiency bleeding (VKDB)) and increased risk of bone fractures. Toxicity of excess vitamin K has not been reported (108, 113).

2.5. Liver Disease and Occurrence of Fat-soluble Vitamin Deficiency

Fat-soluble vitamin deficiency is observed in patients suffering from liver diseases of various etiologies. A recent study reported deficiencies of vitamin A (29%), vitamin D (71%), vitamin E (6%) and up to 68% showed signs of subclinical vitamin K deficiency, in a group of 31 patients suffering from cholestatic liver disease (114). Similarly, a prevalence of fat-soluble vitamin A, D, E and K deficiency is reported to be 33.5%, 13.2%, 1.9% and 7.8%, respectively, in a group of over 150 PBC patients (115). In PSC patients a prevalence of vitamin A, D and E deficiency was found of respectively 40%, 14% and 2%. In pre-transplant patients prevalence was increased to 82%, 57% and 43%, respectively (116, 117). A recent study by De Paula *et al.* reports 60% of patients (35 out of 58) suffering from liver cirrhosis have lowered plasma retinol levels, 23 of these patients were diagnosed as vitamin A deficient (VAD) (118). More than half (54.3% out of $n = 140$) of patients suffering from hepatitis C virus-related chronic liver disease were VAD (119). The vast majority (85%) of patients ($n = 147$) with alcoholic liver cirrhosis have a compromised vitamin D status, 37% of these patients were vitamin D deficient (VDD). In the same study, 47% of PBC patients ($n = 58$), had lowered vitamin D serum levels, 18% of these patients were VDD (120). Sokol *et al.* reported that 17% of PBC patients (7 out of 42) suffered from vitamin E deficiency. In multiple studies, patients suffering from PBC, biliary atresia and autoimmune hepatitis presented lower serum levels of vitamins A, D, E and K when compared to controls, without reaching deficiency (120-123). The prevalence of fat-soluble vitamin deficiencies varies between studies because of differences in disease etiology, group size, patient age and the definition of “deficiency”. Collectively, these studies identify vitamin A deficiency as the most frequent deficiency of all fat-soluble vitamins in (cholestatic) liver diseases.

2.6. Supplementation of Fat-soluble Vitamins and Liver Disease

Guidelines for supplementation of fat-soluble vitamins to patients suffering from liver disease differ per country. In literature, fat-soluble vitamin supplementation is recommended in patients suffering from cholestatic liver disease, especially in neonates and infants, as stores of fat-soluble vitamins are low at birth (124, 125). Vitamin A supplementation is recommended as a single 50.000 IU dose every 15 days for adults and every month for children. However, no clear guidelines exist for vitamin A supplementation for patients with liver disease in The Netherlands. Administration of vitamin E, 100-200-400 IU/day, is recommended for all patients with cholestasis and neurological symptoms of unknown etiology. Vitamin K malabsorption is rapidly corrected by subcutaneous administering of vitamin K, 10 mg/day for 3 days, followed by long-term oral supplements of vitamin K, 5-10 mg/day or 10 mg/month subcutaneously (126, 127).

When fat-soluble vitamin supplements are given attention should be paid to the route of administering fat-soluble vitamin supplements. Oral supplementation of fat-soluble vitamins to (obstructive) cholestatic patients may not be as efficient as subcutaneous supplementation, since fat absorption is disturbed in these patients because of limited bile flow to the intestine.

3. VITAMIN A METABOLISM

3.1. Ileal Vitamin A Absorption Requires Bile Salts

Various forms of -precursors of- vitamin A are entering the digestive tract depending on the composition of the diet. Plant carotenoids are thought to be absorbed into the intestinal epithelium by passive diffusion after being incorporated into micelles that mainly consist of bile salts and dietary fats. However, recent studies in Caco-2 monolayers suggest a facilitated uptake mechanism of carotenoids that involves scavenger receptor class B member 1 (SR-BI) (128, 129). Retinyl esters are first converted to retinol within the intestinal lumen, followed by uptake in the enterocyte (87). Unesterified retinol enters the cell by diffusion, but the efflux of retinol may be partially facilitated by the ABC transporter ABCA1 (128). The pancreatic triglyceride lipase (PTL) and the intestinal brush border membrane enzyme phospholipase B are responsible for the hydrolysis of retinyl esters. The enzymatic activity of PTL depends on the presence of bile. Lipases, such as PTL, act on a water-lipid interface and bile aids in the formation of such interfaces. The emulsifying properties of bile salts stabilize the interface and bile phospholipids are part of the interface (130). Administration of bile salt sequestering agents to humans lowers the total carotenoid levels in serum (131), while administration of the bile salt TCA enhanced vitamin A absorption in rats (132), further underscoring the role of bile salts in vitamin A absorption.

3.2. Vitamin A Transport

Beta-carotene can be converted to retinoids inside the enterocyte (87). Symmetric cleavage of one molecule β -carotene by beta-carotene 15,15'-monooxygenase 1

(BCMO1) yields two molecules of retinal (133). Alternatively, via a series of asymmetric cleavages one molecule of β -carotene can be converted to one molecule of retinal. Subsequently, retinal is reduced in a reversible process to retinol. Several enzymes are capable of catalysing this conversion, including members of the short- and medium-chain alcohol dehydrogenase/reductase superfamily that will be discussed later. The intestinal reductase activity, however, remains to be identified (134).

Free retinol is bound by the cellular retinol-binding protein type II (CRBP-II/RBP2). Most of the retinol absorbed by the enterocyte is re-esterified to saturated long-chain fatty acids, mainly palmitic acid. Binding of retinol to CRBP-II facilitates the esterification of retinol by lecithin:retinol acyl transferase (LRAT). Uncleaved carotenoids and newly-synthesized retinyl esters, but not the ingested retinyl esters, are incorporated into chylomicrons (CMs) and secreted into lymph (128, 135, 136).

Chylomicrons are very heterogeneously-sized particles that consist of a core of triglycerides and cholesterol esters and a monolayer of phospholipids, cholesterol and proteins. CMs are excreted by the enterocyte in the postprandial state after fat ingestion. CM excretion is impaired in the absence of bile salts (137). Two clinical manifestations of impaired CM assembly are abetalipoproteinemia and CM retention disease. Patients suffer from neurologic disorders and visual impairment. In a mouse model for CM retention disease it was observed that the absorption of fat, vitamin A and E was severely impaired and the mice showed significantly reduced growth rates (135), underscoring the importance of CM in the efficient absorption of fat and fat-soluble vitamins, such as vitamin A.

Although most retinoids leave the enterocyte as retinyl esters, also a considerable amount of free retinol is released into the portal circulation (130).

3.3 Vitamin A Storage and Distribution

Chylomicrons distribute nutrients to the various tissues and the chylomicron remnants are subsequently cleared by the hepatocyte. Most of the retinyl esters are still present in these remnants. Within the hepatocyte, retinyl esters are again hydrolysed to retinol. The trafficking of vitamin A throughout the body is depicted in **Figure 1.4**. Free hepatic retinol is now either used directly for distribution to tissues or stored in the liver. For distribution throughout the organism, retinol associates with RBP4, followed by secretion into the blood circulation. Approximately 95 % of plasma retinol-RBP4 is complexed with transthyretin (TTR) in a 1:1 ratio. This interaction reduces glomerular filtration of retinol (88, 138). Extrahepatic tissues take up retinol by means of the RBP4-receptor “Stimulated by Retinoic Acid gene 6 homolog” (STRA6). The retinol-RBP4-TTR complex dissociates at the receptor and free retinol is taken up by the cell. Free RBP4 in the circulation is catabolized in the kidney tubules. The second direction is transport of retinol to, and storage in, the hepatic stellate cell. Here, free retinol is esterified with long chain fatty acids, again predominantly palmitate, for storage in cytosolic lipid droplets. Retinol esters make up 30-50% of the lipid content of the lipid droplets in stellate cells (90).




Figure 1.4. Major pathways for retinoid transport throughout the body

Prior to absorption, dietary retinyl esters (RE) are hydrolysed to retinol (ROH) in the intestinal lumen, while carotenoids are absorbed and converted intracellularly to retinol by the enterocyte. In the enterocyte, retinol reacts with fatty acids to form retinyl esters, which are incorporated into chylomicrons (CM). Via the intestinal lymph chylomicrons reach the general circulation. Chylomicron remnants (CMR), which still contain most of the retinyl esters, are formed within the blood capillaries. These are cleared by the liver parenchymal cells and to some extent by extrahepatic tissues. In the hepatocytes, retinyl esters are hydrolysed to retinol, which then binds to cellular retinol binding protein 1 (CRBP1/RBP1). The retinol-RBP1 complex is secreted and transported into hepatic stellate cells. Retinol entering the stellate cell is either stored in lipid droplets as retinol palmitate or bound by plasma retinol-binding protein 4 (RBP4) and secreted into blood. In the bloodstream most retinol-RBP4 is reversibly complexed with transthyretin (TTR). Uncomplexed retinol is taken up by cells via a receptor mediated mechanism. Free RBP4 in the circulation is catabolized in the kidney tubules. Based on Senoo (2004) (85) and Blomhoff *et al.* (2006) (83).

Although the mechanism for retinol storage in, and mobilization from, stellate cells has not been fully elucidated, the factor shuttling retinol between different liver cell types is the cellular retinol-binding protein (CRBP-I/RBP1). The involvement and recycling of RBP4 (139) has been ruled out, since mice lacking *Rbp4* accumulate excess of vitamin A in the liver. While still able to store vitamin A, these mice were unable to mobilise it to plasma. Moreover, extrahepatic expression of RBP4 does not restore vitamin A mobilisation in these animals, indicating that circulating RBP4 is not re-used by the liver. STRA6 is not expressed in the liver, ruling out a role of this receptor in hepatic storage of retinoids (140, 141).

Under vitamin A sufficient conditions, most of the retinyl esters absorbed from the chylomicron remnants are transferred as retinol to the stellate cells, where over 80%

of the body supply of vitamin A is stored (88). Smaller amounts are also stored in lipid droplets in the hepatocytes and in stellate cells residing in extrahepatic organs and tissues, such as kidneys, digestive tract, adrenal gland, lung, testis, uterus, bone marrow and thymus (90, 142-144). Extrahepatic storage sites of retinyl esters may provide a local supply of vitamin A to tissues with a high demand, such as the retina. The importance of extrahepatic vitamin A pools is demonstrated by the observation that storage of retinyl esters in retinal pigment epithelial cells is prerequisite for normal visual function (144). The stores of vitamin A are sufficient to maintain a steady physiological concentration of 1-2 μM retinol in plasma, despite strong fluctuation in daily intake of vitamin A (144).

3.4. Vitamin A Metabolism

The three main circulating metabolites of vitamin A in the body differ in their oxidation state: the hydroxyl/alcohol form (retinol), the aldehyde form (retinal) and the carboxylic form (retinoic acid, RA) (96) (**Figure 1.5**). As described above, vitamin A is distributed throughout the body as retinol. It is generally believed that conversion of retinol into the biologically active retinal and retinoic acid occurs locally (88, 145). Retinol is converted into retinoic acid in a two-step process (reviewed in (145, 146)). Three enzyme families are capable of catalysing the first step in which retinol is reversibly converted into retinal: the cytosolic medium-chain alcohol dehydrogenases (ADH), the (membrane-associated) short chain dehydrogenases (SDR) and the cytochrome P450s (CYPs) (88, 147-149). In the second step, retinal is converted into RA in an irreversible reaction. Multiple enzymes contain the retinaldehyde dehydrogenase (RALDH) activity. Best characterized are the mouse and human RALDH1, RALDH2 and RALDH3 that have been shown to be involved in RA synthesis *in vivo* (145). Of these three enzymes, only RALDH1 prefers *cis*-retinaldehyde over all-*trans* retinal with the highest activity towards 9-*cis* retinal (150). Also various enzymes belonging to the human cytochrome P-450 family have been shown to be capable of metabolizing retinals to retinoic acids. These include CYP1A1, CYP1A2 and CYP3A4, of which CYP1A2 shows the highest activity for 9-*cis* retinal (151).

The expression levels of the metabolic enzymes involved in RA synthesis may affect the ratio between atRA and the *cis*-RAs, since these enzymes differ in their substrate specificity. *In vivo* isomerization between atRA and 9cRA has been described and may contribute in fine-tuning individual retinoic acid levels (152).

Although produced locally, RA may act everywhere in the organism. The circulation contains low levels of atRA (0.2 to 0.7 % of plasma retinol). The plasma atRA contributes to variable extent to the tissue pool of RA, depending on the tissue/organ. In liver and brain, the retinoic acid pool originates primarily from the circulation rather than from local synthesis (153). RA concentrations in tissue are very low, on average 3-15 μg per kilogram (88). Therefore, its contribution to dietary intake of vitamin A is negligible. Still, it has been shown that animals grow normally on retinoic acid as the sole source of vitamin A-derivatives indicating that RA is the main bioactive derivative of vitamin A (142).




Figure 1.5. Retinol is converted into retinoic acid in a two-step process

The three main pools of vitamin A in the body differ in their oxidation state: the hydroxyl/alcohol form (retinol), the aldehyde form (retinal) and the carboxylic form (retinoic acid). Vitamin A is distributed throughout the body as (RBP4 bound-) retinol. Conversion of retinol into retinoic acid is believed to occur locally within the target cell. In a two-step process, retinol is first reversibly converted into retinal by members of the cytosolic medium-chain alcohol dehydrogenases (ADH), the (membrane-associated) short chain dehydrogenases (SDR) families and various cytochrome P450 enzymes (CYPs). In the second step retinal is irreversibly converted into retinoic acid by retinaldehyde dehydrogenases (RALDHs). Retinoic acid regulates gene transcription via the retinoic acid and retinoid X receptor. Degradation of retinoic acid is facilitated by various CYPs. Beta-carotene is converted to retinal by beta-carotene 15,15'-monooxygenase 1 (BCMO1) and subsequently converted to retinol and stored as retinylesters in the liver. AKP = aldo-keto reductase (AKR) superfamily; LRAT = lecithin:retinol acyl transferase; REH = retinyl ester hydrolase. Based on Pares *et al.* (2008) (143).

Total RA levels are regulated by an interplay between RA synthesizing and catabolizing enzymes. Retinoic acid is metabolized in a two-step process. In the first step, phase I enzymes of the CYP450 superfamily catabolize the different isomers of RA. One of these enzymes is retinoic acid-inactivating cytochrome P450 (P450RAI/CYP26). This metabolizes atRA, but does not appreciably metabolize cis-RA isomers (9-cis and 13-cis). AtRA induces expression of CYP26, thereby controlling its own catabolism (154, 155). While numerous CYP enzymes have been identified that catabolize atRA and/or 13-cis RA, only a few 9cRA-metabolising enzymes have been identified. Of the CYPs that are dominantly expressed in the adult human liver, CYP2C9, -2C8 and -3A4 are the major ones involved in 9-cis RA metabolism. The efficiency of these CYPs to metabolize 9cRA is higher than that for atRA, which may explain why the concentrations of 9cRA *in vivo* are low. In the second step, phase II enzymes facilitate the conjugation of phase I metabolites. All trans-RA and its phase I metabolites 4-oxo-RA, 5,6-epoxy-RA, and 4-OH-RA were found to be glucuronidated by the human glucuronyl transferase UGT2B7 (147, 156).

4. NUCLEAR RECEPTORS/METABOLIC SENSORS IN CONTROL OF BILE SALT HOMEOSTASIS

In the past decade, bile salts have been shown to be key signalling molecules in various biological pathways, in addition to their long-known fat-emulsifying actions. With the discovery of the bile acid-activated Farnesoid X Receptor (FXR) and its

transcriptional regulation of specific target genes, it has become clear how bile acids control their own synthesis and transport. FXR belongs to the superfamily of nuclear receptors (NRs). Many metabolic processes, including glucose, lipid and cholesterol/bile acid metabolism are regulated by NRs. These are ligand-activated transcription factors. Ligands to these NRs are often small lipophilic molecules including hormones, metabolites and food-derived compounds.

4.1 Nuclear Receptor Superfamily

At present, 49 members of the nuclear receptor (NR) superfamily have been identified (reviewed by Francis *et al.* (157)). The superfamily of NRs is divided into seven subfamilies (NR0 to NR6) based on sequence homology. The nomenclature of nuclear receptors has been standardized per subfamily (reviewed by Germain *et al.* (158)). An in depth overview of the seven subclasses and their members is beyond the scope of this review (see review by Aranda *et al.* (159)). The relevant subfamilies for this thesis will be briefly discussed. Members of the NR1 thyroid hormone receptor-like subfamily include the thyroid hormone receptor (TR), retinoic acid receptor (RAR), peroxisome proliferator-activated receptor (PPAR), constitutive androstane receptor (CAR), liver X receptor (LXR), vitamin D receptor (VDR), pregnane X receptor (PXR) and FXR. The NR2 subfamily, or retinoid X receptor-like nuclear receptors, include hepatocyte nuclear factor-4 (HNF-4) and retinoid X receptor-alpha (RXR α). In general, members of the NR1 subfamily need to form a heterodimer with RXR α to become transcriptionally active (160, 161). In several cases, NRs were identified before their ligands were known. These receptors were therefore designated “orphan receptors”. As their natural and synthetic ligands became known, some of these orphan receptors have been “adopted” (162). Two nuclear receptors that remain orphan to date are the liver receptor homolog-1 (LRH-1) belonging to the NR5 subfamily of steroidogenic factor-like receptors and the small heterodimer partner (SHP), belonging to the NR0 family of miscellaneous NRs.

NRs have a modular structure consisting of multiple functional domains. A typical nuclear receptor consists of a variable N-terminal region (region A/B), a conserved DNA-binding domain (DBD) (region C), a linker (region D), and a conserved E region that contains the ligand binding domain (LBD). Some receptors contain also a C-terminal region (F) with unknown function. Isoforms (alpha, beta, gamma) originate from homologous genes and isoforms (alpha 1, alpha 2) of nuclear receptors are generated via alternative translation initiation sites and/or alternative mRNA splicing. The ligand-independent transcriptional activation domain (AF-1) is contained within the A/B region, and the ligand-dependent transactivation domain (AF-2) is located within the C-terminal portion of the LBD (159).

Nuclear receptors regulate gene transcription by binding to specific DNA sequences, so-called hormone responsive elements (HREs), in the promoter region of specific genes. NR/RXR α heterodimers bind to a variety of tandem repeats of the hexamers AGGTCA or AGTTCA typically spaced by 1 or more base pairs (163, 164). The orientation of these hexamers, or so-called half-sites, may vary giving rise to a direct

repeat (DR) (AGGTCAnAGGTCA), a palindromic everted repeat (ER) (TGACCTnAGGTCA) or a palindromic inverted repeat (IR) (AGGTCAnTGACCT) (**Figure 1.6**). It should be noted that these HREs are consensus sequences and that the actual genomic HREs may slightly differ from the consensus.

Gene regulation by NRs is, however, far more complex than just receptor binding to a responsive element in a promoter. Competition between agonists and antagonists, RXR α availability, heterodimerization efficiency, cofactor recruitment and NR protein modification, together determine the ultimate transcriptional efficiency (165).




Figure 1.6. NR1 class nuclear receptors interact as a dimer with RXR α with hormone response element

RXR α is the obligate heterodimer partner of most members of the NR1 thyroid hormone receptor like subfamily, including the thyroid hormone receptor (TR), retinoic acid receptor (RAR), peroxisome proliferator-activated receptor (PPAR), constitutive androstane receptor (CAR), liver X receptor (LXR) pregnane X receptor (PXR), farnesoid X receptor (FXR) and vitamin D receptor (VDR). The NR/RXR α heterodimer interacts with so called hormone responsive elements (HREs), in the promoter region of certain genes. These HREs typically consists of tandem repeat of the hexamers AGGTCA or AGTTCA spaced by 1 or more bases, giving rise to a direct repeat (DR), a palindromic everted repeat (ER) or a palindromic inverted repeat (IR).

Based on Baranowski. (2008) (362).

4.2. Vitamin A Receptors

RXR and RAR, both activated by vitamin A metabolites, were among the first nuclear receptors that were characterized.

4.2.A. Retinoid X Receptor

The retinoid X receptor (RXR) was identified by Mangelsdorf *et al.* in 1990 (167). Three isotypes of RXR exist, -alpha (RXR α /NR2B1), -beta (RXR β /NR2B2) and -gamma (RXR γ /NR2B3). For each isotype two isoforms have been identified: RXR α (α 1 and α 2), RXR β (β 1 and β 2), and RXR γ (γ 1 and γ 2) (168). RXR α is highly expressed in liver and to a lesser extend in lung, muscle, kidney and spleen. RXR γ is expressed in spleen, adrenal gland, heart, intestine and kidney. RXR β is rather ubiquitously expressed, but relatively low in liver and intestine (169).

The natural ligand of RXR is the vitamin A-derivative 9cRA (169-173). Physiological levels of 9cRA have proven difficult to detect in mammalian tissue (153, 174-176), but have been reported (177, 178). This has triggered some scepticism to whether 9cRA is the main endogenous ligand of RXR (179). Other ligands that have been

shown to modulate RXR activity include poly unsaturated fatty acids (PUFAs), such as docosahexaenoic acid, arachidonic acid and oleic acid (180, 181). Even though 9cRA may not be the only endogenous ligand for RXR, various studies showed that vitamin A signalling (or lack of vitamin A signalling) via RXR affects expression of NR/RXR target genes, supporting that vitamin A-derivatives are physiological ligands for RXR (**Chapters 2 and 3** of this thesis).

Homodimeric RXR interacts with DR-1 sequences (182). More importantly though, RXR is the obligate heterodimer partner of most nuclear receptors belonging to the NR1 subfamily. As such, RXR is a key factor in a multitude of metabolic processes, including bile salt homeostasis.

4.2.B. Retinoic Acid Receptor

As for RXR, also three isotypes of the retinoic acid receptor (RAR) exist, -alpha (RAR α /NR1B1), -beta (RAR β /NR1B2) and -gamma (RAR γ /NR1B3) (183). Each isotype has several isoforms, of which the expression has been extensively studied in mouse embryonic development. RAR α is ubiquitously expressed, while RAR β and RAR γ show a more tissue- and cell type-specific distribution. RAR β is present in the liver capsule as well as in the epithelium and outer mesenchyme of the intestine. RAR γ is largely absent from the gastrointestinal tract, with the exception of the squamous epithelium of the stomach (184).

RAR forms heterodimers with RXR (185). RAR/RXR typically interacts with retinoic acid response elements (RAREs) consisting of a direct repeat of AGGTCA interspaced by 5 nucleotides (DR-5) (186). The main natural ligand of RAR is atRA. RARs also bind 9cRA, but with lower affinity than atRA (187, 188). Target genes of RAR are numerous and include genes involved in vitamin A metabolism, such as RBP4 and RBP2 (189, 190).

RAR/RXR target genes involved in bile acid homeostasis include *Ntcp* (191, 192) and *ASBT* (84). Also rat *Cyp7a1* is reported to be an RAR/RXR target gene (193). Furthermore, RAR/RXR represses *MRP3* transcription (194).

4.3. Bile Acid Receptors

Three members of the nuclear receptor family have been identified as bile acid sensors. FXR is regarded the primary bile acid receptor, regulating transcription of key genes involved in bile acid synthesis and transport. The other two bile acid sensing NRs are PXR and VDR. These are mainly involved in the adaptive response to prevent bile acid hepatotoxicity. In addition, CAR is also activated by increased levels of bile salts. Although CAR is not a bile acid sensor by itself, bile salts and bilirubin induces nuclear translocation of CAR, thereby indirectly activating this NR. Like PXR and VDR, CAR is involved in the detoxification of bile salts. These receptors are extensively reviewed by Moore *et al.* 2006 (195). An additional bile acid receptor is the G protein-coupled receptor TGR5. Unlike the intracellular NRs, TGR5 is a cell surface receptor, responding to extracellular bile acids (196, 197). TGR5 signalling is involved in energy expenditure (198) and does not seem to play a role in bile

acid synthesis. Therefore, functions of TGR5 will be not discussed in this chapter. An overview of bile acid receptor-mediated gene expression of genes involved in bile salt synthesis, transport and metabolism is depicted in **Figure 1.7**.




Figure 1.7. Bile sensors promote bile acid transport and detoxification

Bile acid-activated nuclear receptors promote transport, conjugation and detoxification of bile salts. While FXR primarily dictates the cellular efflux of bile salt, PXR and VDR mainly regulate the detoxification of bile salts. Dotted lines represent VDR/PXR signalling in general, meaning not restricted to bile salt-induced signalling.

4.3.A. Farnesoid X Receptor Is the Mammalian Bile Salt Sensor

The farnesoid X receptor (FXR/NR1H4) was originally identified in 1995 as a retinoid receptor-interacting protein (RIP14, (199)) and found to be activated by farnesol (200), retinoic acid and TTNPB (4-[E-2-(5,6,7,8-tetrahydro-5,5,8,8-tetramethyl-2-naphthalenyl)-1-propenyl] benzoic acid) (201). FXR was named after its first identified ligand, although a direct interaction with any of the above compounds was never established.

A few years later, it was established that bile acids are highly potent physiological activators of FXR. The strongest activating bile acid is CDCA (202-204). With the discovery of its endogenous ligands FXR was renamed bile acid receptor (BAR) (202), but the name FXR is still commonly used today. Other endogenous ligands of FXR include androsterone (205), PUFAs (206) and oxysterol 22(R)-hydroxycholesterol (207). Synthetic ligands include GW4064 (208) and 6-Ethyl-CDCA (6-ECDCA) (209). Plant-derived guggulsterone can function as an FXR agonist (210), but also as an FXR antagonist (211).

FXR is profoundly expressed in tissues that are exposed to bile salts, such as liver and intestine, but also in kidney and adrenal glands (200, 212). Four isoforms of FXR have been identified in rodents and humans, designated Fxrα1, Fxrα2, Fxrβ1, and Fxrβ2

(213). To complicate matters, a homolog of *NR1H4* was identified, *NR1H5*, which was named *FXR β* . *FXR β* encodes a functional receptor for lanosterol in rodents and other mammals but is a pseudogene in humans (214, 215). The conserved *FXR α* (generally termed *FXR/NR1H4*) gene encodes four protein products. Because of differential translation initiation sites, *FXR α 3* (formerly “Fxr β 1”) and *FXR α 4* (formerly “Fxr β 2”) differ at their N-terminus. As a result of differential transcript splicing, *FXR α 1* (“Fxr α 1”) and *FXR α 3* contain a four amino acid-insert in the hinge region that is not present in *FXR α 2* (“Fxr α 2”) and *FXR α 4* (195). The *FXR* isoforms are differentially expressed. In adult humans, *FXR α 1/2* mRNA is predominant in liver and adrenal gland. Expression of *FXR α 3/4* mRNA is most abundant in colon, duodenum, and kidney. *FXR α 3/4* mRNA levels are generally lower than that of *FXR α 1/2* (213).

For most transcriptional targets, *FXR* needs to form a heterodimer with *RXR α* to be transcriptionally active (201). The typical *FXR* responsive element (FXRE) is an inverted repeat spaced by one base pair (IR-1), with the consensus sequence AG-GTCAnTGACCT (200). FXREs have been identified in the promoter regions of genes involved in bile acid homeostasis, including *BSEP* (216, 217), *OST α / β* (218, 219), *IBABP* (204, 220), *BAAT/BAT* (221), bile acid-CoA synthetase (*BACS*) (221), *OATP1B3* (222), *SHP* (223–225) and within the second intron of fibroblast growth factor 19 (*FGF19*) in humans (226) and its rodent ortholog *Fgf15* (227). Alternative *FXR* binding sequences were detected in the promoter elements of *MRP2* (an everted repeat spaced by 8 base pairs; ER-8) (228), sulfotransferase 2A1 (*SULT2A1/STD*) (an inverted repeat without spacing; IR-0) (229), UDP glucuronosyltransferase 2B4 (*UGT2B4*) and the insulin-responsive glucose transporter 4 (*GLUT4*) (both a hexamer half-site) (230, 231). In the latter two, *FXR* induces transcription in the absence of *RXR α* . *FXR*-mediated transcriptional suppression has been described for apolipoprotein A-I (*Apo-AI*) and C-III (*Apo-CIII*) (232, 233). An overview of *FXR* target genes is given in **Table 1.1**.

FXR appears to be a multipurpose nuclear receptor (234), as it regulates expression of numerous target genes beyond bile acid synthesis and transport. Target genes of *FXR* are involved in processes such as blood clotting (fibrinogen (*FBG*) (235, 236) and complement factor 3 (*C3*) (227)), inflammation/adhesion (kininogen (*KNG*) (237) and intercellular adhesion molecule 1 (*ICAM-1/CD54*) (238)) and lipid and cholesterol homeostasis (fatty acid synthase (*FAS*) (239), phospholipid transfer protein (*PLTP*) (240) and apolipoprotein C-II (*Apo-CII*) (241)). Furthermore, *FXR* has been associated with liver regeneration and regrowth, (242) with glucose metabolism (reviewed in (243, 244)) and differentiation of bone marrow stromal cells into osteoblasts (245).

Studies with *Fxr*^{-/-} mice revealed the central role of *FXR* in bile salt homeostasis. Wild type mice tolerate CA-feeding by reducing bile acid synthesis (*Cyp7a1*) and hepatic bile salt import (*Ntcp*), while simultaneously increasing bile salt excretion via *BSEP*. This regulation is strongly impaired in *Fxr*^{-/-} mice that maintain bile acid synthesis and import upon CA-feeding with consistently low *Bsep* levels. As a consequence, CA-fed *Fxr*^{-/-} mice develop significant bile acid-induced liver damage (79).

Table 1.1. Bile salt sensor FXR is a multipurpose transcription factor

	Direct target gene of FXR	FXRE	Reference
Transport of bile acids and other bile components	BSEP/ABCB11	IR-1	(216, 217)
	IBABP	IR-1	(204, 220)
	MDR3/ABCB4	IR-1	(246)
	MRP2/ABCB2	ER-8	(228)
	OATP1B3/OATP8	IR-1	(222)
	OST α and OST β	IR-1	(218, 219)
Bile acid metabolism	BAAT	IR-1	(221)
	BACS	IR-1	(221)
	CYP3A4	IR-1/DR-3, ER-8	(247)
	CYP8B1	IR-1	(248)
	SULT2A1 (STD)	IR-0	(229)
	UGT2B4	Half-site	(230)
	UGT2B7	negFXRE	(249)
Negative feedback on bile acid transport/synthesis	FGF19/Fgf15	IR-1	(226, 227)
	SHP	IR-1	(223-225)
Lipid/cholesterol metabolism	Apo-CII	IR-1	(250)
	Apo-CIII	negFXRE (DR-1)	(233)
	Apo-A1	negFXRE	(232)
	FAS	IR-1	(239)
	INSIG-2	IR-1	(251)
	PLTP	IR-1	(240)
	SR-BI	DR-8	(252)
Blood clotting/inflammation/adhesion	C3	IR-1	(227)
	Fibrinogen (FBG)	ND	(235)
	ICAM-1/CD54	IR-1	(238)
	Kininogen (KNG)	IR-1	(237)
Miscellaneous	ALAS1	IR-1	(253)
	alphaA-crystallin (CRYAA)	IR-1	(254)
	ASCT2	IR-1	(255)
	AT2R	IR-2	(256)
	DDAH1	IR-1	(257)
	eNOS	IR-2	(258)
	Fetuin-B	IR-1	(259)
	GLUT4	Half-site	(231)
	NAGS	ER-8	(260)
	NaS-1/Slc13a1	IR-1	(261)
	OAT2	negFXRE (DR-1)	(262)
	PXR	IR-1	(263)
	Syndecan-1	DR-1	(264)
	VPAC-1	IR-1	(265)

The farnesoid X receptor (FXR) is the primary mammalian bile acid receptor. By interacting with FXR, bile salts promote their own transport (e.g. bile salt export pump (BSEP)) and metabolism/clearance (cytochrome P450 3A4 (CYP3A4)). Via a negative feedback loop which involves the FXR target genes *SHP* and *FGF19*, bile acids suppress the synthesis of *de novo* bile acids. However, FXR signaling is not limited to bile salt synthesis and transport. FXR target genes are also involved in processes such as blood clotting (fibrinogen (*FBG*) and complement factor 3 (*C3*)), inflammation/adhesion (kininogen (*KNG*) and intercellular adhesion molecule 1 (*ICAM-1/CD54*)) and lipid and cholesterol homeostasis (fatty acid synthase (*FAS*), phospholipid transfer protein (*PLTP*) and apolipoprotein C-II (*Apo-C2*)) and glucose metabolism (insulin-responsive glucose transporter type 4 (*GLUT4*)).

4.3.B. Pregnane X Receptor

The pregnane X receptor (PXR/NR1I2), also called the steroid and xenobiotic receptor (SXR), is highly expressed in liver and intestine. PXR is activated by xenobiotic compounds, steroids and bile acids, including LCA, CDCA and DCA (266-271).

The PXR/RXR α heterodimer binds to so-called xenobiotic response elements (XREs) present in the promoter region of genes encoding drug metabolizing enzymes and drug transport proteins. These XREs consist of specific repeats of AG(G/T)TCA oriented as either a direct repeat spaced by three to five nucleotides (DR-3, DR-4 or DR-5) or as an everted repeat spaced by six or eight nucleotides (ER-6 and ER-8) (272). PXR is a master regulator of drugs and xenobiotics metabolism (273) and is involved in the protective response to bile salt hepatotoxicity (270). PXR/RXR α induces transcription of *CYP3A4/Cyp3a11* (267) and *SULT2A1* (274) that respectively catalyse hydroxylation and sulfoconjugation of compounds such as drugs, xenobiotics and bile salts. Hydroxylation and sulfoconjugation makes these compounds more hydrophilic (polar, water-soluble), less toxic and more amenable for urinary excretion (82). Some of the membrane-embedded transport proteins that excrete these detoxified compounds are regulated by PXR and also transport (metabolites of) bile salts, including MRP3 (275), MRP2 (228) and MDR1 (273, 276). Expression of the organic anion transporter *OATP1B1*, which imports bile salts, is also regulated by PXR (272). LCA-activated PXR was shown to induce expression of murine *Cyp3a11* and *Atp1a4* (269).

Besides bile salt transport, PXR activity influences bile acid synthesis. Studies with *Pxr*^{-/-} mice showed that PXR represses *Cyp7a1* expression (269), which include both SHP- and FGF15-dependent mechanisms. The human *SHP* promoter contains a PXR/RXR binding site and LCA-activated PXR was shown to induce *FGF19* promoter activity (277). Both SHP and FGF19 (278, 279) repress *CYP7A1* expression (discussed in section 5.3). To make matters more complex, PXR expression itself is regulated by FXR (263). This may further potentiate the role of PXR to bile acid signalling. Thus, bile acid signalling through PXR likely contributes to the adaptive response to prevent bile acid toxicity.

4.3.C. Vitamin D Receptor

In addition to its main natural ligand 1,25-dihydroxy vitamin D₃ (1,25-(OH)₂D₃), the vitamin D receptor (VDR; NR1I1) is also activated to the same extent by LCA and 3-keto-LCA (280, 281). VDR regulates calcium homeostasis and bone metabolism (reviewed by St-Arnaud (282)). The VDR/RXR heterodimer binds to vitamin D response elements (VDREs). The typical VDRE consists of a direct hexameric (GGTCCA, AGGTCA, or GGGTGA) repeat spaced by 3 nucleotides (DR-3), although the number of spacing nucleotides may vary (283). Less frequent, everted repeats have also been found to act as VDREs (reviewed in (284)).

With regard to bile acid metabolism, LCA-activated VDR was shown to induce expression of *CYP3A4* (280, 281, 285), *MDR1* (286) and *MRP3* (287), while *Cyp7a1* expression is repressed by LCA-activated VDR (286). In addition, VDR signalling

induces *SULT2A1* (288, 289) expression and was reported to inhibit the CDCA-dependent transcription of the FXR target genes *SHP*, *IBABP* and *BSEP* (290). VDR thus actively participates in the detoxification and clearance of bile acids, fine-tunes expression of FXR target genes and inhibits bile acid biosynthesis.

4.3.D. Constitutive Androstane Receptor

The constitutive androstane receptor (CAR; NR1I3) is highly expressed in liver and, as the name suggests, is a constitutive active receptor (291, 292). Transcriptional activity of CAR has different modes of action (review by Tzameli *et al.* (293)). Although CAR does not require a ligand to become transcriptionally active (294), ligands have been identified that modulate its constitutive activity. CAR plays a role in bilirubin clearance as is apparent from studies with *Car*^{-/-} mice challenged with a single dose of bilirubin. Bilirubin itself was shown to activate CAR. However, bilirubin does not seem to be a direct agonist of CAR (295), but rather promotes its nuclear translocation. In a similar fashion certain bile acids are able to indirectly activate CAR by inducing nuclear translocation (296). The CAR/RXR heterodimer interacts with RAREs (291). CAR protects against bile acid toxicity in CA-fed *Fxr* and *Pxr* double knockout mice. Pretreatment with CAR activators resulted in decreased bile acid and bilirubin serum levels, through upregulation of hepatic genes involved in bile acid and/or bilirubin metabolism and excretion (297). Similar results were obtained comparing LCA-fed *Car*^{-/-} and WT mice (298). Target genes of CAR involved in bile acid detoxification and clearance include cytosolic sulfotransferases (*SULTs*) (299), 3'-phosphoadenosine 5'-phosphosulfate synthetase 2 (*Pappss2*) (299), *Cyp3a11* (300), *Mrp3* and *Mrp4* (reviewed in (14)).

4.3.E. Additional Nuclear Receptors Involved in Bile Acid Metabolism

Two orphan NRs have a prominent role in the transcriptional regulation of bile acid homeostasis. The liver receptor homolog-1 (LRH-1) induces transcription of various genes involved in bile acid transport and synthesis, while the small heterodimer partner (SHP) is involved in a negative feedback loop, counteracting the activity of LRH-1 and various other NR/RXR heterodimers. The interplay between LRH-1 and SHP is one of the main mechanisms that fine-tunes bile salt synthesis and transport.

LRH-1 induces bile salt synthesis and transport

The liver receptor homolog-1 (LRH-1/NR5A2), also known as fetoprotein transcription factor (FTF), is expressed in liver, intestine, pancreas, ovary and adrenal glands (301-303). It has also been named CYP7A promoter-binding factor, as it is well-known as a positive regulator of expression of hepatic *Cyp7A1* (301). Unlike most NRs, LRH-1 does not form a heterodimer with RXR, but instead binds DNA as a monomer to an extended nuclear receptor half-site YCAAGGYCR (where Y is any pyrimidine and R is any purine) (302, 304).

No endogenous (or synthetic) ligand has been identified to date that is associated with the constitutive activity of LRH-1. Although *in silico* modelling studies pre-

dict that phospholipids may directly bind to LRH-1 (reviewed by Forman (305)), the biological relevance of this putative interaction remains elusive. Murine LRH-1 was shown to exist as a stable monomer in the absence of a ligand, suggesting that it might not need a ligand at all to be transcriptionally active (306).

LRH-1 responsive elements (LRE) have been identified in the promoter regions of *Cyp7a1*, *Cyp8b1*, *ASBT*, *MRP3*, *BSEP*, *OST α /OST β* , *ABCG5/ABCG8* and *SHP* (84, 301, 307-313). LRH-1 facilitates the LXR α -mediated expression of rodent *Cyp7A1* (225, 314) and *Fas* (315) *in vitro*. Hepatic *Lrh-1*^{-/-} mice showed that *Cyp8b1*, but not *Cyp7a1* transcription critically depends on LRH-1 *in vivo* (316, 317). Also murine hepatic and intestinal *Shp* transcription depends largely on LRH-1 (316, 317).

Transactivation by LRH-1 seems to be required for maximal induction of FXR target genes in response to bile salts, as was shown for *BSEP* (311) and *MRP3*. For the latter, two adjacent LREs were identified within its promoter. Mutating either of these responsive elements resulted in loss of CDCA-induced *MRP3* promoter activity (318). Additionally, a recent study demonstrated that LRH-1 recruits FXR to the promoter of genes involved in lipid metabolism (319).

Bile acid-induced transcription of *LRH-1/Lrh-1* is observed in human hepatic and intestinal cell lines and in rat liver (224, 310, 318, 320). However, involvement of FXR in LRH-1 expression has not been established. Interestingly, bile acid-regulated expression of LRH-1 was not observed in mice (225). Taken together, current knowledge establishes LRH-1 as a general positive regulator of bile acid and cholesterol homeostasis and in some species bile salts may increase LRH-1 expression.

Small Heterodimer Partner Is a Repressor of Bile Salt Homeostasis

The small heterodimer partner (SHP/NR0B2) is an atypical member of the NR-family, as it lacks a DNA-binding domain and a natural ligand has not been identified to date. However, recent evidence suggests that SHP activity can be modulated in a ligand-dependent fashion, albeit by synthetic compounds (e.g. 4-[3-(1-adamanty-1)-4-hydroxyphenyl]-3-chlorocinnamic acid (3-Cl-AHPC)) (321). *SHP* is expressed in liver, small intestine, adrenal gland and spleen (322). SHP is a general negative transcription factor of signalling pathways involving nuclear receptors and plays an important role in the negative feedback on bile salt synthesis and transport. However, functions of SHP extend far beyond bile salt homeostasis, including control of viral replication (323), tumor suppression (324), sexual maturation (325), obesity and (type 2) diabetes (326, 327).

Transcription of *SHP/Shp* is induced by FXR (223-225) and LXR α (in humans) (328). As a consequence, *Shp* transcription is low in *FXR*^{-/-} mice and CA-feeding does not alter *Shp* transcription in these animals (79). SHP represses gene transcription by directly interacting with other NRs, cofactors or chromatin-modifying enzymes (322, 329). An important SHP-target involved in positive regulation of bile acid synthesis and transport is *LRH-1*. The role of SHP in the negative feedback of bile salt homeostasis will be discussed in section 5.

Liver X Receptor

Two isoforms of the liver X receptor (LXR) are described; LXR α (NR1H3) and LXR β (NR1H2) (330). *LXR α* is profoundly expressed in liver, intestine, kidney, adipose tissue and spleen (330, 331), while *LXR β* is ubiquitously expressed (332). The typical LXR/RXR responsive element (LXRE) is a hexanucleotide sequence (AGGTCA) separated by four nucleotides (DR-4) (330, 331). LXRs are activated by cholesterol-derivatives and intermediates of cholesterol metabolism, including the oxysterols 22(R)-hydroxycholesterol, and 24(S)-hydroxycholesterol, but also 6- α -hydroxy bile acids (333-336). LXR signalling controls cholesterol, fatty acid and carbohydrate metabolism (reviewed in (337, 338)). *Lxr α ^{-/-}* mice suffer from impaired cholesterol and bile acid metabolism (339), indicating the interconnected regulation of bile acid, cholesterol, and lipid metabolism (162, 302, 340). Target genes of LXR that are involved in bile acid and cholesterol metabolism include *Cyp7a1* (334, 339), *ABCG5* and *ABCG8* (341), *SHP* (328) and *LRH-1* (342). Notably, expression of rodent, but not human *CYP7A1*, is regulated via LXR (328). LXR and FXR share, or compete for, responsive elements in the promoter regions of *hIBABP* and *mOsta α/β* (343, 344). FXR signalling has been suggested to affect cholesterol transport via *ABCG5* and *ABCG8* (345). CA feeding of mice resulted in increased expression of hepatic *Lxr α* , but FXR does not seem to control *Lxr* expression directly (346). LXR signalling thus influences bile acid synthesis and co-regulates transcription of certain FXR-target genes involved in bile salt transport.

4.4. Permissive Versus Nonpermissive Functioning of RXR/9cRA Within the NR/RXR-Heterodimer

In NR/RXR heterodimers, 2 different ligands play a role in modulating the transcriptional activity of the protein complex. The presence of 9cRA as the RXR ligand may have different effects on the NR/RXR activity depending on the specific NR and/or target gene studied and have been subclassified as permissive and nonpermissive NR/RXR complexes (159, 347). In permissive NR/RXR heterodimers, RXR is both a structural and functional component allowing for 9cRA signalling through such a NR/RXR heterodimer. Ligands of RXR and its NR partner can independently and synergistically activate gene transcription. This has been described for PPARs (348, 349), LXR (330) and also FXR (220, 240).

In nonpermissive NHR/RXR heterodimers RXR is merely a structural component of the heterodimer required for DNA-binding, but not necessarily acting as a receptor. Moreover, 9cRA may even repress expression of the target genes (347). RXR is non-permissive in heterodimers with RAR (350), TR (351) and VDR (352, 353). However, a given NR/RXR combination is not strictly permissive or nonpermissive, as this also depends on the specific target gene. For instance, RXR is a permissive partner to TR in prolactin gene regulation (354). While RXR/CAR heterodimers seem to be neither strictly permissive nor nonpermissive, as the contribution of RXR/9cRA depends on the experimental approach. Tzameli *et al.* therefore argue that classifying a particular NR/RXR complex as either permissive or nonpermissive is an oversimplification (355).

The RXR/FXR heterodimer has been described as permissive, based on the regulation of *IBABP* and *PLTP* (220, 240). We and others have shown that RXR α acts as a nonpermissive partner to FXR in the regulation of human and mouse *BSEP* (**Chapter 2**, Kassam *et al.* (356)).

5. FEEDBACK REGULATION ON BILE SALT SYNTHESIS AND TRANSPORT

Increased levels of circulating bile salts repress *de novo* synthesis of bile acids, hepatic and intestinal bile salt uptake and promote bile salt export, detoxification and clearance. Both FXR-dependent and -independent mechanisms have been identified in this process. A schematic overview of this feedback regulation on synthesis and transport is depicted in **Figures 1.8** and **1.9**.

5.1. The FXR-SHP(-LRH-1) Cascade

Bile acid-activated FXR/RXR α enhances the expression of *SHP* (223, 224), which in turn represses the expression of hepatic *CYP7A1* (223, 224), *CYP8B1* (357), *Ntcp* (191) and ileal *ASBT* (84, 358), thereby reducing biosynthesis and cellular import of bile salts.

Many genes that are transcriptionally repressed by SHP are LRH-1 target genes. Interaction of SHP with LRH-1 prevents LRH-1-dependent gene transcription. SHP interacts with the AF-2 domain of LRH-1 and competes with co-activators for binding to this domain. Once bound to its target receptor and recruited to DNA, the inherent repression function of SHP further contributes to the overall inhibitory effect (359, 360).

SHP also represses the LRH-1-enhanced LXR α -dependent transcription of rodent *Cyp7a1* (223, 225). *Cyp7a1* and *Cyp8b1* levels are only slightly elevated in *Shp*^{-/-} mice compared to WT animals (346, 361-363) under normal conditions. However, these animals are more susceptible to bile duct-ligation-induced cholestasis, as *Cyp7a1* (and *Cyp8b1*) transcription is not efficiently repressed in the absence of SHP (363). Transcriptional regulation of rodent *Cyp7a1* differs from human *CYP7A1* as the LXRE is not conserved in the human *CYP7A1* promoter (364, 365). Activated LXR α represses *CYP7A1* transcription in primary human hepatocytes (328). Moreover, human *CYP7A1* is repressed by bile acids and this mechanism may still include SHP, as *CYP7A1* is repressed by SHP in primary human hepatocytes (328) as well. Thus, SHP is of key importance in the adaptive response to increasing hepatic bile acid concentrations.

Expression of hepatic and ileal bile salt import proteins is also repressed in response to rising bile salt concentrations. Expression of rat *Ntcp* is regulated by hepatocyte nuclear factor 1 (HNF1) (366, 367), hepatocyte nuclear factor 4 alpha (HNF4 α) (368) and RAR/RXR (192). SHP represses transcription of rat *Ntcp* by interfering with RAR/RXR-dependent transcription (191) and by lowering the HNF4 α -dependent transcription of *Hnf1 α* (369). Not much is known about the bile acid-dependent repression mechanism on human NTCP expression.




Figure 1.8. Bile salts transport and synthesis is regulated by bile salts themselves

The farnesoid X receptor (FXR) is the primary mammalian bile acid receptor. FXR acts as a heterodimer with the retinoid X receptor (RXR). By interacting with FXR, bile salts promote their own transport (e.g. bile salt export pump (BSEP)) and organic solute transporter α/β (OST α/β). Via a negative feedback loop that involves the FXR target genes small heterodimer partner (SHP) and fibroblast growth factor (FGF19), bile salts suppress the *de novo* synthesis of bile acids. Both SHP and FGF19 inhibit expression of cholesterol 7- α -hydroxylase (CYP7A1). SHP inhibits transcription by interfering with binding of the transcription factor liver receptor homolog-1 (LRH-1) to its responsive element within the promoter region of CYP7A1, while ileal expressed FGF19 interacts with the hepatic fibroblast growth factor receptor 4 (FGFR4). This leads in a multi-step signalling pathway to phosphorylation of c-Jun N-terminal kinase (JNK), which in turn inhibits CYP7A1 expression. Hepatic and ileal uptake of bile salts by the sodium/taurocholate co-transporting polypeptide (NTCP) and ileal bile salt transporter (ASBT), respectively, is repressed by SHP. Both NTCP and ASBT are RAR/RXR α target genes. SHP interferes with the RAR/RXR α -dependent transcription of these genes. Notably, the human liver also expresses FGF19, while expression of the rodent ortholog fibroblast growth factor 15 (FGF15) seems limited to the ileum.

Human and mouse, but not rat ASBT expression is repressed by bile acids. This species difference is explained by the absence of an LRE in the rat *Asbt* promoter. Moreover, LRH-1 seems not to be expressed in rat ileal epithelial cells (84, 309). ASBT expression in *Fxr*^{-/-} mice is unresponsive to bile acid feeding (309). Both LRH-1 and RAR/RXR responsive elements have been identified in the human *ASBT* promoter region. Mutating either of these two sequences was found to reduce *ASBT* promoter activity, but only the RARE appeared to be required for bile acid-dependent inhibiti-

on of *ABST* expression (84). Expression of both human *NTCP* and *ASBT* is strongly increased by the nuclear receptor glucocorticoid receptor (GR/NR3C1). *ASBT/NTCP* activation by GR can be inhibited by SHP (370, 371).

Inhibition of BSEP, OST α/β and MRP3 expression via LRH-1 and SHP does not seem to contribute to the cytoprotective mechanisms activated by increasing intracellular bile acid levels, as it prevents the clearance of bile acids.

5.2. Questioning the Role of LRH-1 In Negative Feedback Regulation

Interestingly, studies with liver- and intestine-specific *Lrh-1* deficient mice showed that LRH-1 is not required for the negative feedback regulation of bile acid synthesis, irrespective of its established role in bile acid homeostasis (316, 317). *Cyp7a1* and *Cyp8b1* transcription is higher in *Lrh-1*^{+/-} heterozygote mice than in WT mice, which is accompanied by an increased bile salt concentration in bile. Overexpression of LRH-1 in mice repressed both *Cyp7a1* and *Cyp8b1* transcription, suggesting that LRH-1 might also act as a repressor of bile acid biosynthesis (372). *Shp*^{-/-} mice have slightly elevated *Cyp7a1*, *Cyp8b1* and *Bsep* mRNA levels (346, 361). The increase in *Bsep* is probably the result of the increased bile acid pool size in these animals (346). However, lack of SHP may also increase *Bsep* expression by lack of inhibition of LRH-1, like observed for *Cyp7a1* and *Cyp8b1* (311). Together, these studies indicate that both the basal levels of the constitutive active LRH-1 and its interaction with SHP determine its effect on bile acid homeostasis.

5.3. Redundant Pathways of Bile Acid Synthesis Repression

In contrast to what would be expected from the feedback control mechanisms described above, *Cyp7a1* expression is strongly increased in bile duct-ligated rodents, despite high concentrations of bile salts in the liver and blood. Feeding these bile duct-ligated mice the FXR ligand GW4064 decreased hepatic *Cyp7a1* expression (279). Also CA feeding of *Shp*^{-/-} mice repressed *Cyp7a1* and *Cyp8b1* transcription (361, 362). These findings stipulate the existence of FXR-SHP-independent feedback mechanisms to repress bile acid biosynthesis.

5.3.A. FXR - FGF19 - JNK Mediated Repression of CYP7A1 Expression

An alternative mode of CYP7A1 repression is provided by *FGF19* (*Fgf15* in rodents) (226). Bile acids from the intestinal lumen activate ileal FXR/RXR α and increase expression of *Fgf15*. Ileal FGF15 is released in the blood and signals to the liver via the hepatic fibroblast growth factor receptor 4 (FGFR4), which leads to phosphorylation of JNK (226, 279). Activated JNK represses CYP7A1 expression (226, 346, 373). Consequently, both *Fgf15*^{-/-} and *Fgfr4*^{-/-} mice display higher *Cyp7a1* mRNA expression than WT mice and an inability to repress *Cyp7a1* in response to feeding of FXR ligands. *Asbt*^{-/-} mice have lower *Fgf15* transcription and higher *Cyp7a1* and *Cyp8b1* transcription levels than WT mice (374), indicating that reabsorbed bile acids in the ileum determine the rate of *de novo* bile acid biosynthesis.




Figure 1.9. Bile acid sensors engage in negative feedback on bile acid biosynthesis and cellular import

Bile salt-activated FXR suppresses bile acid synthesis (CYP7A1) via SHP and FGF19. Inhibition of CYP7A1 via FGF19 involves the phosphorylation of JNK. SHP also limits cellular uptake of bile salts by inhibiting NTCP and ASBT expression. In addition to FXR signalling, bile acid signalling via PXR and VDR also contribute to the negative feedback regulation on bile acid synthesis. PXR signalling promotes expression of FGF19 and SHP. Furthermore, interaction between the ligand-activated PXR and SHP proteins enhances inhibition of CYP7A1 (dotted line). Moreover, PXR is an FXR-target gene itself. Bile acid signalling via VDR inhibits CYP7A1 by competing with other transcription factors at the *CYP7A1* promoter. FXR-independent CYP7A1 repression mechanisms may converge the FXR-dependent mechanism on JNK phosphorylation.

Additionally, FGF15/FGF19 was recently shown to inhibit ASBT expression (375), expending FXR-FGF15/FGF19 signalling to bile acid transport. Thus, the FXR-SHP and the FXR-FGF19 signalling cascades are both involved in feedback mechanisms that control bile salt synthesis and transport.

5.3.B. SHP and FGF19 Pathways Overlap

SHP-independent mechanisms of CYP7A1 repression seem to dominate over SHP dependent mechanisms, as *Cyp7a1* repression prevails in the absence of *Shp* (362), but not in the absence of *Fgf15* (279). However, SHP does not seem to be fully dispensable in the negative feedback loop of bile acid biosynthesis. Activation of either FXR or RXR by feeding mice synthetic ligands GW4064 or LG100268, respectively, was found to repress *Cyp7A1* transcription, but this was dependent on SHP (346,

361). However, CA-feeding reduced *Cyp7a1* (and *Cyp8b1*) mRNA levels in *Shp*^{-/-} mice, but this seems to depend on hepatotoxicity rather than on FXR signalling (361). Moreover, the FGF15- and SHP-dependent repression of *Cyp7a1* overlap, since SHP significantly enhances the FGF15 mediated repression of *Cyp7a1* mRNA (279).

5.3.C. FXR Independent Repression

The negative feedback loop on bile acid biosynthesis also includes FXR-independent mechanisms. LCA-activated PXR (269) or VDR (376) represses *Cyp7a1* gene expression. This mode of repression may involve FGF19/FGF15, as it has been shown that the *FGF19* promoter is activated by LCA-activated PXR in intestinal cells (277). Alternatively, LCA-activated PXR is suggested to bind to the bile acid response element (BARE)-I in the *CYP7A1* promoter and inhibits *CYP7A1* transcription by both SHP-dependent and -independent pathways. PXR strongly interacts with SHP in a ligand-dependent manner (377).

FXR-SHP-independent bile acid-induced expression of inflammatory cytokines (e.g., tumor necrosis factor alpha (TNF α) and interleukin beta (IL-1 β)) by Kupffer cells may mediate *CYP7A1* repression as well (378, 379). Ito and co-workers reported that mice lacking the membrane protein beta-klotho (KLB) have increased *Cyp7a1* and *Cyp8b1* mRNA expression, while bile acid-dependent repression of these genes was still largely intact. Recently, KLB has been shown to modulate FGF19/FGF15 signalling via FGFR4 (380, 381). This links the observation that KLB inhibits ASBT expression, to the repression of ASBT by FGF19/FGF15 (375).

5.3.D. FXR Dependent and Independent Repression of CYP7A1 Converges at JNK

Bile acid signalling via FGF15/19, TNF α , IL-1 β and protein kinase C (PKC) (377, 382) all activate JNK. Activated JNK in turn activates jun proto-oncogene (JUN), which inhibits the HNF4 α -dependent transcription of *CYP7A1* (383), and induces *Shp* expression (373). Additionally, phosphorylation of FXR by PKC promotes its transcriptional activity (384). This way, signalling via both FXR dependent and independent pathways may converge and amplify each other in the effort to repress *CYP7A1* expression.

6. AIM AND OUTLINE OF THIS THESIS

Vitamin A and bile acid homeostasis are interrelated at various levels. Bile salts are required for hydrolysis of retinyl esters in the intestinal lumen and the absorption of vitamin A into the intestinal epithelium. Consequently, patients suffering from obstructive cholestasis suffer from malabsorption of vitamin A. Synthesis of bile acids takes place in the liver, which also serves as the primary site of vitamin A storage in the mammalian body. Liver disease may progress to liver fibrosis and affect the hepatic stellate cells that contain the vitamin A in intracellular lipid droplets. During fibrosis, stellate cells become activated and differentiate to proliferative and contractile myofibroblasts that overproduce extracellular matrix proteins leading to the characteristic scar tissue formation. In this process they lose their retinoid content. Thus cholestasis effectively promotes depletion of vitamin A from the organism by emptying the stores and preventing uptake.

Bile acids and vitamin A-derivatives like 9-*cis* retinoic acid (9cRA) co-regulate bile acid synthesis in the liver and transport between the liver and intestine. They act mainly by activating transcription factors of the nuclear receptor family, in particular the farnesoid X receptor (FXR) and the retinoid X receptor- α (RXR α). FXR is activated by bile acids and RXR α by 9cRA and they act together as a heterodimer in regulating expression of genes involved in bile acid synthesis and transport.

The role of bile acid-activated FXR in regulating bile acid homeostasis has been well established. However, the role of its heterodimer partner RXR remains largely illusive. As many chronic liver diseases may develop to vitamin A deficiency, the putative role of vitamin A in bile salt homeostasis is therefore highly relevant.

In this thesis, we aim to determine the role of vitamin A in the FXR/RXR α -mediated regulation of bile acid homeostasis both *in vitro* and *in vivo* models.

In **chapter 2**, we performed *in vitro* and *in vivo* experiments to analyse the effect of vitamin A on the bile acid-induced expression of the transporter that is responsible for secretion of bile salts from the liver, the bile salt export pump (BSEP).

In **chapter 3**, we analysed the effect of vitamin A on the bile acid-induced expression of the small heterodimer partner (SHP), which is a transcriptional repressor of several genes in bile acid synthesis and transport. Even though expression of both BSEP and SHP is induced by FXR, we observed opposite effects of 9cRA on their expression, repressing BSEP and super-inducing SHP. **Chapter 2** and **3** include a detailed analysis of the molecular mechanisms that cause this opposite effect of 9cRA on BSEP and SHP expression.

Given the fact that vitamin A-derivatives indeed directly affect expression of genes involved in bile salt biosynthesis and transport, we analysed in **chapter 4** whether vitamin A deficiency in laboratory animals leads to changes in bile salt homeostasis. Finally, in **chapter 5** we analysed whether the combination of vitamin A deficiency and obstructive cholestasis in laboratory animals leads to uncontrolled bile salt regulation and putatively to aggravation of liver damage.

In **chapter 6** an integrated discussion of all our data is presented.

REFERENCES

1. Esteller A. Physiology of bile secretion. *World J Gastroenterol* 2008 Oct 7;14(37):5641-5649.
2. Frevert U. Sneaking in through the back entrance: the biology of malaria liver stages. *Trends Parasitol* 2004 Sep;20(9):417-424.
3. Ridlon JM, Kang DJ, Hylemon PB. Bile salt biotransformations by human intestinal bacteria. *J Lipid Res* 2006 Feb;47(2):241-259.
4. Pellicoro A, Faber KN. Review article: The function and regulation of proteins involved in bile salt biosynthesis and transport. *Aliment Pharmacol Ther* 2007 Dec;26 Suppl 2:149-160.
5. Shefer S, Hauser S, Bekersky I, Mosbach EH. Biochemical site of regulation of bile acid biosynthesis in the rat. *J Lipid Res* 1970 Sep;11(5):404-411.
6. Russell DW. The enzymes, regulation, and genetics of bile acid synthesis. *Annu Rev Biochem* 2003;72:137-174.
7. Pellicoro A, van den Heuvel FA, Geuken M, Moshage H, Jansen PL, Faber KN. Human and rat bile acid-CoA:amino acid N-acyltransferase are liver-specific peroxisomal enzymes: implications for intracellular bile salt transport. *Hepatology* 2007 Feb;45(2):340-348.
8. Matysik S, Martin J, Bala M, Scherer M, Schaffler A, Schmitz G. Bile acid signaling after an oral glucose tolerance test. *Chem Phys Lipids* 2011 Sep;164(6):525-529.
9. Li T, Franci JM, Boehme S, Ochoa A, Zhang Y, Klaassen CD, *et al.* Glucose and insulin induction of bile acid synthesis: mechanisms and implication in diabetes and obesity. *J Biol Chem* 2012 Jan 13;287(3):1861-1873.
10. Alrefai WA, Gill RK. Bile acid transporters: structure, function, regulation and pathophysiological implications. *Pharm Res* 2007 Oct;24(10):1803-1823.
11. Kullak-Ublick GA, Stieger B, Meier PJ. Enterohepatic bile salt transporters in normal physiology and liver disease. *Gastroenterology* 2004 Jan;126(1):322-342.
12. Trauner M, Boyer JL. Bile salt transporters: molecular characterization, function, and regulation. *Physiol Rev* 2003 Apr;83(2):633-671.
13. Meier PJ, Stieger B. Bile salt transporters. *Annu Rev Physiol* 2002;64:635-661.
14. Zollner G, Marschall HU, Wagner M, Trauner M. Role of nuclear receptors in the adaptive response to bile acids and cholestasis: pathogenetic and therapeutic considerations. *Mol Pharm* 2006 May;3(3):231-251.
15. Hagenbuch B, Meier PJ. Molecular cloning, chromosomal localization, and functional characterization of a human liver Na⁺/bile acid cotransporter. *J Clin Invest* 1994 Mar;93(3):1326-1331.
16. Hagenbuch B, Stieger B, Foguet M, Lubbert H, Meier PJ. Functional expression cloning and characterization of the hepatocyte Na⁺/bile acid cotransport system. *Proc Natl Acad Sci U S A* 1991 Dec 1;88(23):10629-10633.
17. Meier PJ. Molecular mechanisms of hepatic bile salt transport from sinusoidal blood into bile. *Am J Physiol* 1995 Dec;269(6 Pt 1):G801-G812.
18. Csanaky IL, Lu H, Zhang Y, Ogura K, Choudhuri S, Klaassen CD. Organic anion-transporting polypeptide 1b2 (Oatp1b2) is important for the hepatic uptake of unconjugated bile acids: Studies in Oatp1b2-null mice. *Hepatology* 2011 Jan;53(1):272-281.
19. Hagenbuch B, Dawson P. The sodium bile salt cotransport family SLC10. *Pflugers Arch* 2004 Feb;447(5):566-570.
20. Geyer J, Wilke T, Petzinger E. The solute carrier family SLC10: more than a family of bile acid transporters regarding function and phylogenetic relationships. *Naunyn Schmiedeberg Arch Pharmacol* 2006 Mar;372(6):413-431.
21. Schroeder A, Eckhardt U, Stieger B, Tynes R, Schteingart CD, Hofmann AF, *et al.* Substrate specificity of the rat liver Na⁺-bile salt cotransporter in *Xenopus laevis* oocytes and in CHO cells. *Am J Physiol* 1998 Feb;274(2 Pt 1):G370-G375.

22. Kim JY, Kim KH, Lee JA, Namkung W, Sun AQ, Ananthanarayanan M, *et al.* Transporter-mediated bile acid uptake causes Ca^{2+} -dependent cell death in rat pancreatic acinar cells. *Gastroenterology* 2002 Jun;122(7):1941-1953.
23. Konig J, Seithel A, Gradhand U, Fromm MF. Pharmacogenomics of human OATP transporters. *Naunyn Schmiedebergs Arch Pharmacol* 2006 Mar;372(6):432-443.
24. Konig J, Cui Y, Nies AT, Keppler D. A novel human organic anion transporting polypeptide localized to the basolateral hepatocyte membrane. *Am J Physiol Gastrointest Liver Physiol* 2000 Jan;278(1):G156-G164.
25. Konig J, Cui Y, Nies AT, Keppler D. Localization and genomic organization of a new hepatocellular organic anion transporting polypeptide. *J Biol Chem* 2000 Jul 28;275(30):23161-23168.
26. Kullak-Ublick GA, Hagenbuch B, Stieger B, Scheingart CD, Hofmann AF, Wolkoff AW, *et al.* Molecular and functional characterization of an organic anion transporting polypeptide cloned from human liver. *Gastroenterology* 1995 Oct;109(4):1274-1282.
27. Jacquemin E, Hagenbuch B, Stieger B, Wolkoff AW, Meier PJ. Expression cloning of a rat liver Na^{+} -independent organic anion transporter. *Proc Natl Acad Sci U S A* 1994 Jan 4;91(1):133-137.
28. Foucaud L, Grillasas J, Niot I, Domingo N, Lafont H, Planells R, *et al.* Output of liver fatty acid-binding protein (L-FABP) in bile. *Biochim Biophys Acta* 1999 Jan 4;1436(3):593-599.
29. Monaco HL. Review: the liver bile acid-binding proteins. *Biopolymers* 2009 Dec;91(12):1196-1202.
30. Martin GG, Atshaves BP, McIntosh AL, Mackie JT, Kier AB, Schroeder F. Liver fatty-acid-binding protein (L-FABP) gene ablation alters liver bile acid metabolism in male mice. *Biochem J* 2005 Nov 1;391(Pt 3):549-560.
31. Rembacz KP, Woudenberg J, Hoekstra M, Jonkers EZ, van den Heuvel FA, Buist-Homan M, *et al.* Unconjugated bile salts shuttle through hepatocyte peroxisomes for taurine conjugation. *Hepatology* 2010 Dec;52(6):2167-2176.
32. Suchy FJ, Ananthanarayanan M. Bile salt excretory pump: biology and pathobiology. *J Pediatr Gastroenterol Nutr* 2006 Jul;43 Suppl 1:S10-S16.
33. Gerloff T, Stieger B, Hagenbuch B, Madon J, Landmann L, Roth J, *et al.* The sister of P-glycoprotein represents the canalicular bile salt export pump of mammalian liver. *J Biol Chem* 1998 Apr 17;273(16):10046-10050.
34. van Mil SW, van der Woerd WL, van der BG, Sturm E, Jansen PL, Bull LN, *et al.* Benign recurrent intrahepatic cholestasis type 2 is caused by mutations in ABCB11. *Gastroenterology* 2004 Aug;127(2):379-384.
35. Strautnieks SS, Bull LN, Knisely AS, Kocoshis SA, Dahl N, Arnell H, *et al.* A gene encoding a liver-specific ABC transporter is mutated in progressive familial intrahepatic cholestasis. *Nat Genet* 1998 Nov;20(3):233-238.
36. Jansen PL, Strautnieks SS, Jacquemin E, Hadchouel M, Sokal EM, Hooiveld GJ, *et al.* Hepatocanalicular bile salt export pump deficiency in patients with progressive familial intrahepatic cholestasis. *Gastroenterology* 1999 Dec;117(6):1370-1379.
37. Dean M, Rzhetsky A, Allikmets R. The human ATP-binding cassette (ABC) transporter superfamily. *Genome Res* 2001 Jul;11(7):1156-1166.
38. Akita H, Suzuki H, Ito K, Kinoshita S, Sato N, Takikawa H, *et al.* Characterization of bile acid transport mediated by multidrug resistance associated protein 2 and bile salt export pump. *Biochim Biophys Acta* 2001 Mar 9;1511(1):7-16.
39. Toyoda Y, Hagiya Y, Adachi T, Hoshijima K, Kuo MT, Ishikawa T. MRP class of human ATP binding cassette (ABC) transporters: historical background and new research directions. *Xenobiotica* 2008 Jul;38(7-8):833-862.
40. Kool M, de HM, Scheffer GL, Scheper RJ, van Eijk MJ, Juijn JA, *et al.* Analysis of expression of cMOAT (MRP2), MRP3, MRP4, and MRP5, homologues of the multidrug resistance-associated protein gene (MRP1), in human cancer cell lines. *Cancer Res* 1997 Aug 15;57(16):3537-3547.
41. Gerk PM, Li W, Megaraj V, Vore M. Human multidrug resistance protein 2 transports the therapeutic bile salt tauroursodeoxycholate. *J Pharmacol Exp Ther* 2007 Feb;320(2):893-899.

42. Dubin IN, Johnson FB. Chronic idiopathic jaundice with unidentified pigment in liver cells; a new clinico-pathologic entity with a report of 12 cases. *Medicine (Baltimore)* 1954 Sep;33(3):155-197.
43. Kartenbeck J, Leuschner U, Mayer R, Keppler D. Absence of the canalicular isoform of the MRP gene-encoded conjugate export pump from the hepatocytes in Dubin-Johnson syndrome. *Hepatology* 1996 May;23(5):1061-1066.
44. Paulusma CC, Kool M, Bosma PJ, Scheffer GL, ter BF, Scheper RJ, *et al.* A mutation in the human canalicular multispecific organic anion transporter gene causes the Dubin-Johnson syndrome. *Hepatology* 1997 Jun;25(6):1539-1542.
45. Kuipers F, Enserink M, Havinga R, van der Steen AB, Hardonk MJ, Fevery J, *et al.* Separate transport systems for biliary secretion of sulfated and unsulfated bile acids in the rat. *J Clin Invest* 1988 May;81(5):1593-1599.
46. Crawford AR, Smith AJ, Hatch VC, Oude Elferink RP, Borst P, Crawford JM. Hepatic secretion of phospholipid vesicles in the mouse critically depends on mdr2 or MDR3 P-glycoprotein expression. Visualization by electron microscopy. *J Clin Invest* 1997 Nov 15;100(10):2562-2567.
47. de Vree JM, Jacquemin E, Sturm E, Cresteil D, Bosma PJ, Aten J, *et al.* Mutations in the MDR3 gene cause progressive familial intrahepatic cholestasis. *Proc Natl Acad Sci U S A* 1998 Jan 6;95(1):282-287.
48. Oude Elferink RP, Ottenhoff R, van WM, Smit JJ, Schinkel AH, Groen AK. Regulation of biliary lipid secretion by mdr2 P-glycoprotein in the mouse. *J Clin Invest* 1995 Jan;95(1):31-38.
49. Paulusma CC, Elferink RP, Jansen PL. Progressive familial intrahepatic cholestasis type 1. *Semin Liver Dis* 2010 May;30(2):117-124.
50. Tauscher A, Kuver R. ABCG5 and ABCG8 are expressed in gallbladder epithelial cells. *Biochem Biophys Res Commun* 2003 Aug 8;307(4):1021-1028.
51. Graf GA, Yu L, Li WP, Gerard R, Tuma PL, Cohen JC, *et al.* ABCG5 and ABCG8 are obligate heterodimers for protein trafficking and biliary cholesterol excretion. *J Biol Chem* 2003 Nov 28;278(48):48275-48282.
52. Yu L, Hammer RE, Li-Hawkins J, Von BK, Lutjohann D, Cohen JC, *et al.* Disruption of Abcg5 and Abcg8 in mice reveals their crucial role in biliary cholesterol secretion. *Proc Natl Acad Sci U S A* 2002 Dec 10;99(25):16237-16242.
53. Thiebaut F, Tsuruo T, Hamada H, Gottesman MM, Pastan I, Willingham MC. Cellular localization of the multidrug-resistance gene product P-glycoprotein in normal human tissues. *Proc Natl Acad Sci U S A* 1987 Nov;84(21):7735-7738.
54. Elias E, Mills CO. Coordinated defence and the liver. *Clin Med* 2007 Apr;7(2):180-184.
55. Stieger B, Fattinger K, Madon J, Kullak-Ublick GA, Meier PJ. Drug- and estrogen-induced cholestasis through inhibition of the hepatocellular bile salt export pump (Bsep) of rat liver. *Gastroenterology* 2000 Feb;118(2):422-430.
56. Wang R, Salem M, Yousef IM, Tuchweber B, Lam P, Childs SJ, *et al.* Targeted inactivation of sister of P-glycoprotein gene (spgp) in mice results in nonprogressive but persistent intrahepatic cholestasis. *Proc Natl Acad Sci U S A* 2001 Feb 13;98(4):2011-2016.
57. Lam P, Wang R, Ling V. Bile acid transport in sister of P-glycoprotein (ABCB11) knockout mice. *Biochemistry* 2005 Sep 20;44(37):12598-12605.
58. Wong MH, Oelkers P, Craddock AL, Dawson PA. Expression cloning and characterization of the hamster ileal sodium-dependent bile acid transporter. *J Biol Chem* 1994 Jan 14;269(2):1340-1347.
59. Wong MH, Oelkers P, Dawson PA. Identification of a mutation in the ileal sodium-dependent bile acid transporter gene that abolishes transport activity. *J Biol Chem* 1995 Nov 10;270(45):27228-27234.
60. Lazaridis KN, Pham L, Tietz P, Marinelli RA, deGroen PC, Levine S, *et al.* Rat cholangiocytes absorb bile acids at their apical domain via the ileal sodium-dependent bile acid transporter. *J Clin Invest* 1997 Dec 1;100(11):2714-2721.
61. Oelkers P, Kirby LC, Heubi JE, Dawson PA. Primary bile acid malabsorption caused by mutations in the ileal sodium-dependent bile acid transporter gene (SLC10A2). *J Clin Invest* 1997 Apr 15;99(8):1880-1887.

62. Kok T, Hulzebos CV, Wolters H, Havinga R, Agellon LB, Stellaard F, *et al.* Enterohepatic circulation of bile salts in farnesoid X receptor-deficient mice: efficient intestinal bile salt absorption in the absence of ileal bile acid-binding protein. *J Biol Chem* 2003 Oct 24;278(43):41930-41937.
63. Praslickova D, Torchia EC, Sugiyama MG, Magrane EJ, Zwicker BL, Kolodzieyski L, *et al.* The ileal lipid binding protein is required for efficient absorption and transport of bile acids in the distal portion of the murine small intestine. *PLoS ONE* 2012;7(12):e50810.
64. Xia X, Francis H, Glaser S, Alpini G, LeSage G. Bile acid interactions with cholangiocytes. *World J Gastroenterol* 2006 Jun 14;12(22):3553-3563.
65. Alpini G, Glaser SS, Rodgers R, Phinizz J, Robertson WE, Lasater J, *et al.* Functional expression of the apical Na⁺-dependent bile acid transporter in large but not small rat cholangiocytes. *Gastroenterology* 1997 Nov;113(5):1734-1740.
66. Lazaridis KN, Tietz P, Wu T, Kip S, Dawson PA, LaRusso NF. Alternative splicing of the rat sodium/bile acid transporter changes its cellular localization and transport properties. *Proc Natl Acad Sci U S A* 2000 Sep 26;97(20):11092-11097.
67. Hirohashi T, Suzuki H, Takikawa H, Sugiyama Y. ATP-dependent transport of bile salts by rat multidrug resistance-associated protein 3 (Mrp3). *J Biol Chem* 2000 Jan 28;275(4):2905-2910.
68. Scheffer GL, Kool M, de HM, de Vree JM, Pijnenborg AC, Bosman DK, *et al.* Tissue distribution and induction of human multidrug resistant protein 3. *Lab Invest* 2002 Feb;82(2):193-201.
69. Zelcer N, van de WK, de WR, Scheffer GL, Marschall HU, Wielinga PR, *et al.* Mice lacking Mrp3 (Abcc3) have normal bile salt transport, but altered hepatic transport of endogenous glucuronides. *J Hepatol* 2006 Apr;44(4):768-775.
70. Rost D, Mahner S, Sugiyama Y, Stremmel W. Expression and localization of the multidrug resistance-associated protein 3 in rat small and large intestine. *Am J Physiol Gastrointest Liver Physiol* 2002 Apr;282(4):G720-G726.
71. Kool M, van der LM, de HM, Scheffer GL, de Vree JM, Smith AJ, *et al.* MRP3, an organic anion transporter able to transport anti-cancer drugs. *Proc Natl Acad Sci U S A* 1999 Jun 8;96(12):6914-6919.
72. Gerk PM, Vore M. The Ileocyte Basolateral Organic Solute Transporter (OST α -OST β) Complex: Finding The Missing Link in Enterohepatic Circulation. *Mol Interv* 2005 Feb;5(1):8-10.
73. Dawson PA, Hubbert M, Haywood J, Craddock AL, Zerangue N, Christian WV, *et al.* The heteromeric organic solute transporter alpha-beta, Ost α -Ost β , is an ileal basolateral bile acid transporter. *J Biol Chem* 2005 Feb 25;280(8):6960-6968.
74. Wang W, Seward DJ, Li L, Boyer JL, Ballatori N. Expression cloning of two genes that together mediate organic solute and steroid transport in the liver of a marine vertebrate. *Proc Natl Acad Sci U S A* 2001 Jul 31;98(16):9431-9436.
75. Seward DJ, Koh AS, Boyer JL, Ballatori N. Functional complementation between a novel mammalian polygenic transport complex and an evolutionarily ancient organic solute transporter, OST α -OST β . *J Biol Chem* 2003 Jul 25;278(30):27473-27482.
76. Ballatori N, Christian WV, Lee JY, Dawson PA, Soroka CJ, Boyer JL, *et al.* OST α -OST β : a major basolateral bile acid and steroid transporter in human intestinal, renal, and biliary epithelia. *Hepatology* 2005 Dec;42(6):1270-1279.
77. Rao A, Haywood J, Craddock AL, Belinsky MG, Kruh GD, Dawson PA. The organic solute transporter alpha-beta, Ost α -Ost β , is essential for intestinal bile acid transport and homeostasis. *Proc Natl Acad Sci U S A* 2008 Mar 11;105(10):3891-3896.
78. Ballatori N. Pleiotropic functions of the organic solute transporter Ost α -Ost β . *Dig Dis* 2011;29(1):13-17.
79. Sinal CJ, Tohkin M, Miyata M, Ward JM, Lambert G, Gonzalez FJ. Targeted disruption of the nuclear receptor FXR/BAR impairs bile acid and lipid homeostasis. *Cell* 2000 Sep 15;102(6):731-744.
80. Ros JE, Libbrecht L, Geuken M, Jansen PL, Roskams TA. High expression of MDR1, MRP1, and MRP3 in the hepatic progenitor cell compartment and hepatocytes in severe human liver disease. *J Pathol* 2003 Aug;200(5):553-560.

81. Boyer JL, Trauner M, Mennone A, Soroka CJ, Cai SY, Moustafa T, *et al.* Upregulation of a basolateral FXR-dependent bile acid efflux transporter OST α -OST β in cholestasis in humans and rodents. *Am J Physiol Gastrointest Liver Physiol* 2006 Jun;290(6):G1124-G1130.
82. Zollner G, Wagner M, Fickert P, Silbert D, Gumhold J, Zatloukal K, *et al.* Expression of bile acid synthesis and detoxification enzymes and the alternative bile acid efflux pump MRP4 in patients with primary biliary cirrhosis. *Liver Int* 2007 Sep;27(7):920-929.
83. Mennone A, Soroka CJ, Cai SY, Harry K, Adachi M, Hagey L, *et al.* MRP4^{-/-} mice have an impaired cytoprotective response in obstructive cholestasis. *Hepatology* 2006 May;43(5):1013-1021.
84. Neimark E, Chen F, Li X, Shneider BL. Bile acid-induced negative feedback regulation of the human ileal bile acid transporter. *Hepatology* 2004 Jul;40(1):149-156.
85. Wolf G. A history of vitamin A and retinoids. *FASEB J* 1996 Jul;10(9):1102-1107.
86. Nomenclature of retinoids: recommendations 1981. IUPAC-IUB Joint Commission on Biochemical Nomenclature (JCBN). *Arch Biochem Biophys* 1983 Jul 15;224(2):728-731.
87. Moore T. Vitamin A and carotene: The absence of the liver oil vitamin A from carotene. VI. The conversion of carotene to vitamin A *in vivo*. *Biochem J* 1930;24(3):692-702.
88. Blomhoff R, Blomhoff HK. Overview of retinoid metabolism and function. *J Neurobiol* 2006 Jun;66(7):606-630.
89. Weber D, Grune T. The contribution of beta-carotene to vitamin A supply of humans. *Mol Nutr Food Res* 2012 Feb;56(2):251-258.
90. Senoo H. Structure and function of hepatic stellate cells. *Med Electron Microsc* 2004 Mar;37(1):3-15.
91. Hix LM, Lockwood SF, Bertram JS. Bioactive carotenoids: potent antioxidants and regulators of gene expression. *Redox Rep* 2004;9(4):181-191.
92. Paiva SA, Russell RM. Beta-carotene and other carotenoids as antioxidants. *J Am Coll Nutr* 1999 Oct;18(5):426-433.
93. Samokyszyn VM, Marnett LJ. Inhibition of liver microsomal lipid peroxidation by 13-cis-retinoic acid. *Free Radic Biol Med* 1990;8(5):491-496.
94. Livrea MA, Tesoriere L, Bongiorno A, Pintaudi AM, Ciaccio M, Riccio A. Contribution of vitamin A to the oxidation resistance of human low density lipoproteins. *Free Radic Biol Med* 1995 Mar;18(3):401-409.
95. Ritter E, Elgeti M, Bartl FJ. Activity switches of rhodopsin. *Photochem Photobiol* 2008 Jul;84(4):911-920.
96. Bonet ML, Ribot J, Felipe F, Palou A. Vitamin A and the regulation of fat reserves. *Cell Mol Life Sci* 2003 Jul;60(7):1311-1321.
97. Holick MF. Vitamin D deficiency. *N Engl J Med* 2007 Jul 19;357(3):266-281.
98. Dusso AS, Brown AJ, Slatopolsky E. Vitamin D. *Am J Physiol Renal Physiol* 2005 Jul;289(1):F8-28.
99. Brown AJ, Dusso A, Slatopolsky E. Vitamin D. *Am J Physiol* 1999 Aug;277(2 Pt 2):F157-F175.
100. Lawson DE, Douglas J, Lean M, Sedrani S. Estimation of vitamin D₃ and 25-hydroxyvitamin D₃ in muscle and adipose tissue of rats and man. *Clin Chim Acta* 1986 Jun 15;157(2):175-181.
101. IUPAC-IUB Joint Commission on Biochemical Nomenclature (JCBN). Nomenclature of tocopherols and related compounds. Recommendations 1981. *Eur J Biochem* 1982 Apr;123(3):473-475.
102. Evans HM, Bishop KS. On the existence of a hitherto unrecognized dietary factor essential for reproduction. *Science* 1922 Dec 8;56(1458):650-651.
103. Assem M, Schuetz EG, Leggas M, Sun D, Yasuda K, Reid G, *et al.* Interactions between hepatic MRP4 and Sult2a as revealed by the constitutive androstane receptor and MRP4 knockout mice. *J Biol Chem* 2004 May 21;279(21):22250-22257.
104. Zingg JM, Azzi A. Non-antioxidant activities of vitamin E. *Curr Med Chem* 2004 May;11(9):1113-1133.
105. Traber MG, Atkinson J. Vitamin E, antioxidant and nothing more. *Free Radic Biol Med* 2007 Jul 1;43(1):4-15.

106. Rigotti A. Absorption, transport, and tissue delivery of vitamin E. *Mol Aspects Med* 2007 Oct;28(5-6):423-436.
107. Kayata S, Kindberg C, Greer FR, Suttie JW. Vitamin K1 and K2 in infant human liver. *J Pediatr Gastroenterol Nutr* 1989 Apr;8(3):304-307.
108. Greer FR. Vitamin K the basics-What's new? *Early Hum Dev* 2010 Jul;86 Suppl 1:43-7.
109. Bolton-Smith C, Price RJ, Fenton ST, Harrington DJ, Shearer MJ. Compilation of a provisional UK database for the phyloquinone (vitamin K1) content of foods. *Br J Nutr* 2000 Apr;83(4):389-399.
110. Dam H. The antihæmorrhagic vitamin of the chick. *Biochem J* 1935 Jun;29(6):1273-1285.
111. Vermeer C, Jie KS, Knapen MH. Role of vitamin K in bone metabolism. *Annu Rev Nutr* 1995;15:1-22.
112. Van WM, De BR, Van d, V, Van BS. Vitamin K, an update for the paediatrician. *Eur J Pediatr* 2009 Feb;168(2):127-134.
113. Fairfield KM, Fletcher RH. Vitamins for chronic disease prevention in adults: scientific review. *JAMA* 2002 Jun 19;287(23):3116-3126.
114. Strole J, Lovell G, Heubi J. Prevalence of subclinical vitamin K deficiency in cholestatic liver disease. *J Pediatr Gastroenterol Nutr* 2009 Jul;49(1):78-84.
115. Phillips JR, Angulo P, Petterson T, Lindor KD. Fat-soluble vitamin levels in patients with primary biliary cirrhosis. *Am J Gastroenterol* 2001 Sep;96(9):2745-2750.
116. Jorgensen RA, Lindor KD, Sartin JS, LaRusso NF, Wiesner RH. Serum lipid and fat-soluble vitamin levels in primary sclerosing cholangitis. *J Clin Gastroenterol* 1995 Apr;20(3):215-219.
117. Sokol RJ, Balistreri WF, Hoofnagle JH, Jones EA. Vitamin E deficiency in adults with chronic liver disease. *Am J Clin Nutr* 1985 Jan;41(1):66-72.
118. de Paula TP, Ramalho A, Braulio VB. The effectiveness of relative dose response to retinol intake as an evaluation of vitamin A status of cirrhotic patients. *J Hum Nutr Diet* 2010 Dec;23(6):583-589.
119. Peres WA, Chaves GV, Goncalves JC, Ramalho A, Coelho HS. Vitamin A deficiency in patients with hepatitis C virus-related chronic liver disease. *Br J Nutr* 2011 Jun 8;1-8.
120. Malham M, Jorgensen SP, Ott P, Agnholt J, Vilstrup H, Borre M, *et al.* Vitamin D deficiency in cirrhosis relates to liver dysfunction rather than aetiology. *World J Gastroenterol* 2011 Feb 21;17(7):922-925.
121. Saron ML, Godoy HT, Hessel G. Nutritional status of patients with biliary atresia and autoimmune hepatitis related to serum levels of vitamins A, D and E. *Arq Gastroenterol* 2009 Jan;46(1):62-68.
122. Floreani A, Baragiotta A, Martines D, Naccarato R, D'odorico A. Plasma antioxidant levels in chronic cholestatic liver diseases. *Aliment Pharmacol Ther* 2000 Mar;14(3):353-358.
123. Kowdley KV, Emond MJ, Sadowski JA, Kaplan MM. Plasma vitamin K1 level is decreased in primary biliary cirrhosis. *Am J Gastroenterol* 1997 Nov;92(11):2059-2061.
124. EASL Clinical Practice Guidelines: management of cholestatic liver diseases. *J Hepatol* 2009 Aug;51(2):237-267.
125. Chapman R, Fevery J, Kalloo A, Nagorney DM, Boberg KM, Shneider B, *et al.* Diagnosis and management of primary sclerosing cholangitis. *Hepatology* 2010 Feb;51(2):660-678.
126. Perez FT, Lopez SP, Tomas E, Gutierrez ML, Lledo JL, Cacho G, *et al.* Diagnostic and therapeutic approach to cholestatic liver disease. *Rev Esp Enferm Dig* 2004 Jan;96(1):60-73.
127. Amedee-Manesme O, Mourey MS, Therasse J, Couturier M, Alvarez F, Hanck A, *et al.* Short- and long-term vitamin A treatment in children with cholestasis. *Am J Clin Nutr* 1988 Apr;47(4):690-693.
128. During A, Harrison EH. Mechanisms of provitamin A (carotenoid) and vitamin A (retinol) transport into and out of intestinal Caco-2 cells. *J Lipid Res* 2007 Oct;48(10):2283-2294.
129. During A, Harrison EH. Intestinal absorption and metabolism of carotenoids: insights from cell culture. *Arch Biochem Biophys* 2004 Oct 1;430(1):77-88.
130. Harrison EH. Mechanisms of digestion and absorption of dietary vitamin A. *Annu Rev Nutr* 2005;25:87-103.

131. Probstfield JL, Lin TL, Peters J, Hunninghake DB. Carotenoids and vitamin A: the effect of hypocholesterolemic agents on serum levels. *Metabolism* 1985 Jan;34(1):88-91.
132. Hollander D. Retinol lymphatic and portal transport: influence of pH, bile, and fatty acids. *Am J Physiol* 1980 Sep;239(3):G210-G214.
133. Von Lintig J, Wyss A. Molecular analysis of vitamin A formation: cloning and characterization of beta-carotene 15,15'-dioxygenases. *Arch Biochem Biophys* 2001 Jan 1;385(1):47-52.
134. Li E, Tso P. Vitamin A uptake from foods. *Curr Opin Lipidol* 2003 Jun;14(3):241-247.
135. Hussain MM. A proposed model for the assembly of chylomicrons. *Atherosclerosis* 2000 Jan;148(1):1-15.
136. Huang HS, Goodman DS. Vitamin A and carotenoids. I. Intestinal absorption and metabolism of 14c-labelled vitamin A alcohol and beta-carotene in the rat. *J Biol Chem* 1965 Jul;240:2839-2844.
137. Davidson NO, Kollmer ME, Glickman RM. Apolipoprotein B synthesis in rat small intestine: regulation by dietary triglyceride and biliary lipid. *J Lipid Res* 1986 Jan;27(1):30-39.
138. Peterson PA. Studies on the interaction between prealbumin, retinol-binding protein, and vitamin A. *J Biol Chem* 1971 Jan 10;246(1):44-49.
139. Norum KR, Blomhoff R. McCollum Award Lecture, 1992: vitamin A absorption, transport, cellular uptake, and storage. *Am J Clin Nutr* 1992 Oct;56(4):735-744.
140. Quadro L, Blaner WS, Hamberger L, Novikoff PM, Vogel S, Piantedosi R, *et al.* The role of extrahepatic retinol binding protein in the mobilization of retinoid stores. *J Lipid Res* 2004 Nov;45(11):1975-1982.
141. Quadro L, Blaner WS, Salchow DJ, Vogel S, Piantedosi R, Gouras P, *et al.* Impaired retinal function and vitamin A availability in mice lacking retinol-binding protein. *EMBO J* 1999 Sep 1;18(17):4633-4644.
142. Erickson JM, Mawson AR. Possible role of endogenous retinoid (Vitamin A) toxicity in the pathophysiology of primary biliary cirrhosis. *J Theor Biol* 2000 Sep 7;206(1):47-54.
143. Nagy NE, Holven KB, Roos N, Senoo H, Kojima N, Norum KR, *et al.* Storage of vitamin A in extrahepatic stellate cells in normal rats. *J Lipid Res* 1997 Apr;38(4):645-658.
144. Blomhoff R, Green MH, Berg T, Norum KR. Transport and storage of vitamin A. *Science* 1990 Oct 19;250(4979):399-404.
145. Duester G, Mic FA, Molotkov A. Cytosolic retinoid dehydrogenases govern ubiquitous metabolism of retinol to retinaldehyde followed by tissue-specific metabolism to retinoic acid. *Chem Biol Interact* 2003 Feb 1;143-144:201-210.
146. Duester G. Families of retinoid dehydrogenases regulating vitamin A function: production of visual pigment and retinoic acid. *Eur J Biochem* 2000 Jul;267(14):4315-4324.
147. Marill J, Idres N, Capron CC, Nguyen E, Chabot GG. Retinoic acid metabolism and mechanism of action: a review. *Curr Drug Metab* 2003 Feb;4(1):1-10.
148. Pares X, Farres J, Kedishvili N, Duester G. Medium- and short-chain dehydrogenase/reductase gene and protein families : Medium-chain and short-chain dehydrogenases/reductases in retinoid metabolism. *Cell Mol Life Sci* 2008 Dec;65(24):3936-3949.
149. Chen H, Howald WN, Juchau MR. Biosynthesis of all-trans-retinoic acid from all-trans-retinol: catalysis of all-trans-retinol oxidation by human P-450 cytochromes. *Drug Metab Dispos* 2000 Mar;28(3):315-322.
150. Labrecque J, Dumas F, Lacroix A, Bhat PV. A novel isoenzyme of aldehyde dehydrogenase specifically involved in the biosynthesis of 9-cis and all-trans retinoic acid. *Biochem J* 1995 Jan 15;305 (Pt 2):681-684.
151. Zhang QY, Dunbar D, Kaminsky L. Human cytochrome P-450 metabolism of retinals to retinoic acids. *Drug Metab Dispos* 2000 Mar;28(3):292-297.
152. Kojima R, Fujimori T, Kiyota N, Toriya Y, Fukuda T, Ohashi T, *et al.* *In vivo* isomerization of retinoic acids. Rapid isomer exchange and gene expression. *J Biol Chem* 1994 Dec 23;269(51):32700-32707.
153. Kurlandsky SB, Gamble MV, Ramakrishnan R, Blaner WS. Plasma delivery of retinoic acid to tissues in the rat. *J Biol Chem* 1995 Jul 28;270(30):17850-17857.
154. Petkovich PM. Retinoic acid metabolism. *J Am Acad Dermatol* 2001 Nov;45(5):S136-S142.

155. Idres N, Marill J, Chabot GG. Regulation of CYP26A1 expression by selective RAR and RXR agonists in human NB4 promyelocytic leukemia cells. *Biochem Pharmacol* 2005 Jun 1;69(11):1595-1601.
156. Marill J, Capron CC, Idres N, Chabot GG. Human cytochrome P450s involved in the metabolism of 9-cis- and 13-cis-retinoic acids. *Biochem Pharmacol* 2002 Mar 1;63(5):933-943.
157. Francis GA, Fayard E, Picard F, Auwerx J. Nuclear receptors and the control of metabolism. *Annu Rev Physiol* 2003;65:261-311.
158. Germain P, Staels B, Dacquet C, Spedding M, Laudet V. Overview of nomenclature of nuclear receptors. *Pharmacol Rev* 2006 Dec;58(4):685-704.
159. Aranda A, Pascual A. Nuclear hormone receptors and gene expression. *Physiol Rev* 2001 Jul;81(3):1269-1304.
160. Zhang XK, Hoffmann B, Tran PB, Graupner G, Pfahl M. Retinoid X receptor is an auxiliary protein for thyroid hormone and retinoic acid receptors. *Nature* 1992 Jan 30;355(6359):441-446.
161. Dong D, Noy N. Heterodimer formation by retinoid X receptor: regulation by ligands and by the receptor's self-association properties. *Biochemistry* 1998 Jul 28;37(30):10691-10700.
162. Chawla A, Repa JJ, Evans RM, Mangelsdorf DJ. Nuclear receptors and lipid physiology: opening the X-files. *Science* 2001 Nov 30;294(5548):1866-1870.
163. Beato M, Herrlich P, Schutz G. Steroid hormone receptors: many actors in search of a plot. *Cell* 1995 Dec 15;83(6):851-857.
164. Forman BM, Evans RM. Nuclear hormone receptors activate direct, inverted, and everted repeats. *Ann N Y Acad Sci* 1995 Jun 12;761:29-37.
165. Karpen SJ. Nuclear receptor regulation of hepatic function. *J Hepatol* 2002 Jun;36(6):832-850.
166. Baranowski M. Biological role of liver X receptors. *J Physiol Pharmacol* 2008 Dec;59 Suppl 7:31-55.
167. Mangelsdorf DJ, Ong ES, Dyck JA, Evans RM. Nuclear receptor that identifies a novel retinoic acid response pathway. *Nature* 1990 May 17;345(6272):224-229.
168. Germain P, Chambon P, Eichele G, Evans RM, Lazar MA, Leid M, *et al.* International Union of Pharmacology. LXIII. Retinoid X receptors. *Pharmacol Rev* 2006 Dec;58(4):760-772.
169. Mangelsdorf DJ, Borgmeyer U, Heyman RA, Zhou JY, Ong ES, Oro AE, *et al.* Characterization of three RXR genes that mediate the action of 9-cis retinoic acid. *Genes Dev* 1992 Mar;6(3):329-344.
170. Heyman RA, Mangelsdorf DJ, Dyck JA, Stein RB, Eichele G, Evans RM, *et al.* 9-cis retinoic acid is a high affinity ligand for the retinoid X receptor. *Cell* 1992 Jan 24;68(2):397-406.
171. Levin AA, Sturzenbecker LJ, Kazmer S, Bosakowski T, Huselton C, Allenby G, *et al.* 9-cis retinoic acid stereoisomer binds and activates the nuclear receptor RXR alpha. *Nature* 1992 Jan 23;355(6358):359-361.
172. Tate BF, Levin AA, Grippo JF. The discovery of 9-cis retinoic acid a hormone that binds the retinoid-X receptor. *Trends Endocrinol Metab* 1994 Jul;5(5):189-194.
173. Allenby G, Bocquel MT, Saunders M, Kazmer S, Speck J, Rosenberger M, *et al.* Retinoic acid receptors and retinoid X receptors: interactions with endogenous retinoic acids. *Proc Natl Acad Sci U S A* 1993 Jan 1;90(1):30-34.
174. Ulven SM, Gundersen TE, Sakhi AK, Glover JC, Blomhoff R. Quantitative axial profiles of retinoic acid in the embryonic mouse spinal cord: 9-cis retinoic acid only detected after all-trans-retinoic acid levels are super-elevated experimentally. *Dev Dyn* 2001 Nov;222(3):341-353.
175. Werner EA, DeLuca HF. Metabolism of a physiological amount of all-trans-retinol in the vitamin A-deficient rat. *Arch Biochem Biophys* 2001 Sep 15;393(2):262-270.
176. Ulven SM, Gundersen TE, Weedon MS, Landaas VO, Sakhi AK, Fromm SH, *et al.* Identification of endogenous retinoids, enzymes, binding proteins, and receptors during early postimplantation development in mouse: important role of retinal dehydrogenase type 2 in synthesis of all-trans-retinoic acid. *Dev Biol* 2000 Apr 15;220(2):379-391.

177. Arnhold T, Tzimas G, Wittfoht W, Plonait S, Nau H. Identification of 9-cis-retinoic acid, 9,13-di-cis-retinoic acid, and 14-hydroxy-4,14-retro-retinol in human plasma after liver consumption. *Life Sci* 1996;59(12):L169-L177.
178. Kane MA, Folias AE, Pingitore A, Perri M, Obrochta KM, Krois CR, *et al.* Identification of 9-cis-retinoic acid as a pancreas-specific autacid that attenuates glucose-stimulated insulin secretion. *Proc Natl Acad Sci U S A* 2010 Dec 14;107(50):21884-21889.
179. Wolf G. Is 9-cis-retinoic acid the endogenous ligand for the retinoic acid-X receptor? *Nutr Rev* 2006 Dec;64(12):532-538.
180. de Urquiza AM, Liu S, Sjoberg M, Zetterstrom RH, Griffiths W, Sjoval J, *et al.* Docosahexaenoic acid, a ligand for the retinoid X receptor in mouse brain. *Science* 2000 Dec 15;290(5499):2140-2144.
181. Lengqvist J, Mata De UA, Bergman AC, Willson TM, Sjoval J, Perlmann T, *et al.* Polyunsaturated fatty acids including docosahexaenoic and arachidonic acid bind to the retinoid X receptor alpha ligand-binding domain. *Mol Cell Proteomics* 2004 Jul;3(7):692-703.
182. Zhang XK, Lehmann J, Hoffmann B, Dawson MI, Cameron J, Graupner G, *et al.* Homodimer formation of retinoid X receptor induced by 9-cis retinoic acid. *Nature* 1992 Aug 13;358(6387):587-591.
183. Germain P, Chambon P, Eichele G, Evans RM, Lazar MA, Leid M, *et al.* International Union of Pharmacology. LX. Retinoic acid receptors. *Pharmacol Rev* 2006 Dec;58(4):712-725.
184. Dolle P. Developmental expression of retinoic acid receptors (RARs). *Nucl Recept Signal* 2009;7:e006.
185. Kliewer SA, Umesono K, Mangelsdorf DJ, Evans RM. Retinoid X receptor interacts with nuclear receptors in retinoic acid, thyroid hormone and vitamin D3 signalling. *Nature* 1992 Jan 30;355(6359):446-449.
186. Umesono K, Murakami KK, Thompson CC, Evans RM. Direct repeats as selective response elements for the thyroid hormone, retinoic acid, and vitamin D3 receptors. *Cell* 1991 Jun 28;65(7):1255-1266.
187. Allenby G, Janocha R, Kazmer S, Speck J, Grippo JF, Levin AA. Binding of 9-cis-retinoic acid and all-trans-retinoic acid to retinoic acid receptors alpha, beta, and gamma. Retinoic acid receptor gamma binds all-trans-retinoic acid preferentially over 9-cis-retinoic acid. *J Biol Chem* 1994 Jun 17;269(24):16689-16695.
188. Repa JJ, Hanson KK, Clagett-Dame M. All-trans-retinol is a ligand for the retinoic acid receptors. *Proc Natl Acad Sci U S A* 1993 Aug 1;90(15):7293-7297.
189. Panariello L, Quadro L, Trematerra S, Colantuoni V. Identification of a novel retinoic acid response element in the promoter region of the retinol-binding protein gene. *J Biol Chem* 1996 Oct 11;271(41):25524-25532.
190. Mangelsdorf DJ, Umesono K, Kliewer SA, Borgmeyer U, Ong ES, Evans RM. A direct repeat in the cellular retinol-binding protein type II gene confers differential regulation by RXR and RAR. *Cell* 1991 Aug 9;66(3):555-561.
191. Denson LA, Sturm E, Echevarria W, Zimmerman TL, Makishima M, Mangelsdorf DJ, *et al.* The orphan nuclear receptor, shp, mediates bile acid-induced inhibition of the rat bile acid transporter, ntcp. *Gastroenterology* 2001 Jul;121(1):140-147.
192. Denson LA, Auld KL, Schiek DS, McClure MH, Mangelsdorf DJ, Karpen SJ. Interleukin-1 β suppresses retinoid transactivation of two hepatic transporter genes involved in bile formation. *J Biol Chem* 2000 Mar 24;275(12):8835-8843.
193. Crestani M, Sadeghpour A, Stroup D, Galli G, Chiang JY. Transcriptional activation of the cholesterol 7 α -hydroxylase gene (CYP7A) by nuclear hormone receptors. *J Lipid Res* 1998 Nov;39(11):2192-2200.
194. Chen W, Cai SY, Xu S, Denson LA, Soroka CJ, Boyer JL. Nuclear receptors RXR α :RAR α are repressors for human MRP3 expression. *Am J Physiol Gastrointest Liver Physiol* 2007 May;292(5):G1221-G1227.
195. Moore DD, Kato S, Xie W, Mangelsdorf DJ, Schmidt DR, Xiao R, *et al.* International Union of Pharmacology. LXII. The NR1H and NR1I receptors: constitutive androstane receptor, pregnane X receptor, farnesoid X receptor alpha, farnesoid X receptor beta, liver X receptor alpha, liver X receptor beta, and vitamin D receptor. *Pharmacol Rev* 2006 Dec;58(4):742-759.
196. Kawamata Y, Fujii R, Hosoya M, Harada M, Yoshida H, Miwa M, *et al.* A G protein-coupled receptor responsive to bile acids. *J Biol Chem* 2003 Mar 14;278(11):9435-9440.

197. Maruyama T, Miyamoto Y, Nakamura T, Tamai Y, Okada H, Sugiyama E, *et al.* Identification of membrane-type receptor for bile acids (M-BAR). *Biochem Biophys Res Commun* 2002 Nov 15;298(5):714-719.
198. Watanabe M, Houten SM, Matakai C, Christoffolete MA, Kim BW, Sato H, *et al.* Bile acids induce energy expenditure by promoting intracellular thyroid hormone activation. *Nature* 2006 Jan 26;439(7075):484-489.
199. Seol W, Choi HS, Moore DD. Isolation of proteins that interact specifically with the retinoid X receptor: two novel orphan receptors. *Mol Endocrinol* 1995 Jan;9(1):72-85.
200. Forman BM, Goode E, Chen J, Oro AE, Bradley DJ, Perlmann T, *et al.* Identification of a nuclear receptor that is activated by farnesol metabolites. *Cell* 1995 Jun 2;81(5):687-693.
201. Zavacki AM, Lehmann JM, Seol W, Willson TM, Kliewer SA, Moore DD. Activation of the orphan receptor RIP14 by retinoids. *Proc Natl Acad Sci U S A* 1997 Jul 22;94(15):7909-7914.
202. Wang H, Chen J, Hollister K, Sowers LC, Forman BM. Endogenous bile acids are ligands for the nuclear receptor FXR/BAR. *Mol Cell* 1999 May;3(5):543-553.
203. Parks DJ, Blanchard SG, Bledsoe RK, Chandra G, Consler TG, Kliewer SA, *et al.* Bile acids: natural ligands for an orphan nuclear receptor. *Science* 1999 May 21;284(5418):1365-1368.
204. Makishima M, Okamoto AY, Repa JJ, Tu H, Learned RM, Luk A, *et al.* Identification of a nuclear receptor for bile acids. *Science* 1999 May 21;284(5418):1362-1365.
205. Wang S, Lai K, Moy FJ, Bhat A, Hartman HB, Evans MJ. The nuclear hormone receptor farnesoid X receptor (FXR) is activated by androsterone. *Endocrinology* 2006 Sep;147(9):4025-4033.
206. Zhao A, Yu J, Lew JL, Huang L, Wright SD, Cui J. Polyunsaturated fatty acids are FXR ligands and differentially regulate expression of FXR targets. *DNA Cell Biol* 2004 Aug;23(8):519-526.
207. Deng R, Yang D, Yang J, Yan B. Oxysterol 22(R)-hydroxycholesterol induces the expression of the bile salt export pump through nuclear receptor farnesoid X receptor but not liver X receptor. *J Pharmacol Exp Ther* 2006 Apr;317(1):317-325.
208. Maloney PR, Parks DJ, Haffner CD, Fivush AM, Chandra G, Plunket KD, *et al.* Identification of a chemical tool for the orphan nuclear receptor FXR. *J Med Chem* 2000 Aug 10;43(16):2971-2974.
209. Pellicciari R, Fiorucci S, Camaioni E, Clerici C, Costantino G, Maloney PR, *et al.* 6 α -ethyl-chenodeoxycholic acid (6-ECDCA), a potent and selective FXR agonist endowed with anticholestatic activity. *J Med Chem* 2002 Aug 15;45(17):3569-3572.
210. Cui J, Huang L, Zhao A, Lew JL, Yu J, Sahoo S, *et al.* Guggulsterone is a farnesoid X receptor antagonist in coactivator association assays but acts to enhance transcription of bile salt export pump. *J Biol Chem* 2003 Mar 21;278(12):10214-10220.
211. Urizar NL, Liverman AB, Dodds DT, Silva FV, Ordentlich P, Yan Y, *et al.* A natural product that lowers cholesterol as an antagonist ligand for FXR. *Science* 2002 May 31;296(5573):1703-1706.
212. Zhang Y, Kast-Woelbern HR, Edwards PA. Natural structural variants of the nuclear receptor farnesoid X receptor affect transcriptional activation. *J Biol Chem* 2003 Jan 3;278(1):104-110.
213. Huber RM, Murphy K, Miao B, Link JR, Cunningham MR, Rupar MJ, *et al.* Generation of multiple farnesoid-X-receptor isoforms through the use of alternative promoters. *Gene* 2002 May 15;290(1-2):35-43.
214. Otte K, Kranz H, Kober I, Thompson P, Hoefer M, Haubold B, *et al.* Identification of farnesoid X receptor beta as a novel mammalian nuclear receptor sensing lanosterol. *Mol Cell Biol* 2003 Feb;23(3):864-872.
215. Zhang ZD, Cayting P, Weinstock G, Gerstein M. Analysis of nuclear receptor pseudogenes in vertebrates: how the silent tell their stories. *Mol Biol Evol* 2008 Jan;25(1):131-143.
216. Plass JR, Mol O, Heegsma J, Geuken M, Faber KN, Jansen PL, *et al.* Farnesoid X receptor and bile salts are involved in transcriptional regulation of the gene encoding the human bile salt export pump. *Hepatology* 2002 Mar;35(3):589-596.
217. Ananthanarayanan M, Balasubramanian N, Makishima M, Mangelsdorf DJ, Suchy FJ. Human bile salt export pump promoter is transactivated by the farnesoid X receptor/bile acid receptor. *J Biol Chem* 2001 Aug 3;276(31):28857-28865.

218. Landrier JF, Eloranta JJ, Vavricka SR, Kullak-Ublick GA. The nuclear receptor for bile acids, FXR, transactivates human organic solute transporter- α and - β genes. *Am J Physiol Gastrointest Liver Physiol* 2006 Mar;290(3):G476-G485.
219. Lee H, Zhang Y, Lee FY, Nelson SF, Gonzalez FJ, Edwards PA. FXR regulates organic solute transporters α and β in the adrenal gland, kidney, and intestine. *J Lipid Res* 2006 Jan;47(1):201-214.
220. Grober J, Zaghini I, Fujii H, Jones SA, Kliewer SA, Willson TM, *et al.* Identification of a bile acid-responsive element in the human ileal bile acid-binding protein gene. Involvement of the farnesoid X receptor/9-*cis*-retinoic acid receptor heterodimer. *J Biol Chem* 1999 Oct 15;274(42):29749-29754.
221. Pircher PC, Kitto JL, Petrowski ML, Tangirala RK, Bischoff ED, Schulman IG, *et al.* Farnesoid X receptor regulates bile acid-amino acid conjugation. *J Biol Chem* 2003 Jul 25;278(30):27703-27711.
222. Jung D, Podvynec M, Meyer UA, Mangelsdorf DJ, Fried M, Meier PJ, *et al.* Human organic anion transporting polypeptide 8 promoter is transactivated by the farnesoid X receptor/bile acid receptor. *Gastroenterology* 2002 Jun;122(7):1954-1966.
223. Goodwin B, Jones SA, Price RR, Watson MA, McKee DD, Moore LB, *et al.* A regulatory cascade of the nuclear receptors FXR, SHP-1, and LRH-1 represses bile acid biosynthesis. *Mol Cell* 2000 Sep;6(3):517-526.
224. Chen W, Owsley E, Yang Y, Stroup D, Chiang JY. Nuclear receptor-mediated repression of human cholesterol 7 α -hydroxylase gene transcription by bile acids. *J Lipid Res* 2001 Sep;42(9):1402-1412.
225. Lu TT, Makishima M, Repa JJ, Schoonjans K, Kerr TA, Auwerx J, *et al.* Molecular basis for feedback regulation of bile acid synthesis by nuclear receptors. *Mol Cell* 2000 Sep;6(3):507-515.
226. Holt JA, Luo G, Billin AN, Bisi J, McNeill YY, Kozarsky KF, *et al.* Definition of a novel growth factor-dependent signal cascade for the suppression of bile acid biosynthesis. *Genes Dev* 2003 Jul 1;17(13):1581-1591.
227. Li J, Pircher PC, Schulman IG, Westin SK. Regulation of complement C3 expression by the bile acid receptor FXR. *J Biol Chem* 2005 Mar 4;280(9):7427-7434.
228. Kast HR, Goodwin B, Tarr PT, Jones SA, Anisfeld AM, Stoltz CM, *et al.* Regulation of multidrug resistance-associated protein 2 (ABCC2) by the nuclear receptors pregnane X receptor, farnesoid X-activated receptor, and constitutive androstane receptor. *J Biol Chem* 2002 Jan 25;277(4):2908-2915.
229. Song CS, Echchgadda I, Baek BS, Ahn SC, Oh T, Roy AK, *et al.* Dehydroepiandrosterone sulfotransferase gene induction by bile acid activated farnesoid X receptor. *J Biol Chem* 2001 Nov 9;276(45):42549-42556.
230. Barbier O, Torra IP, Sirvent A, Claudel T, Blanquart C, Duran-Sandoval D, *et al.* FXR induces the UGT2B4 enzyme in hepatocytes: a potential mechanism of negative feedback control of FXR activity. *Gastroenterology* 2003 Jun;124(7):1926-1940.
231. Shen H, Zhang Y, Ding H, Wang X, Chen L, Jiang H, *et al.* Farnesoid X receptor induces GLUT4 expression through FXR response element in the GLUT4 promoter. *Cell Physiol Biochem* 2008;22(1-4):1-14.
232. Claudel T, Sturm E, Duez H, Torra IP, Sirvent A, Kosykh V, *et al.* Bile acid-activated nuclear receptor FXR suppresses apolipoprotein A-I transcription via a negative FXR response element. *J Clin Invest* 2002 Apr;109(7):961-971.
233. Claudel T, Inoue Y, Barbier O, Duran-Sandoval D, Kosykh V, Fruchart J, *et al.* Farnesoid X receptor agonists suppress hepatic apolipoprotein CIII expression. *Gastroenterology* 2003 Aug;125(2):544-555.
234. Lee FY, Lee H, Hubbert ML, Edwards PA, Zhang Y. FXR, a multipurpose nuclear receptor. *Trends Biochem Sci* 2006 Oct;31(10):572-580.
235. Anisfeld AM, Kast-Woelbern HR, Lee H, Zhang Y, Lee FY, Edwards PA. Activation of the nuclear receptor FXR induces fibrinogen expression: a new role for bile acid signaling. *J Lipid Res* 2005 Mar;46(3):458-68.
236. Plass JRM. Function and regulation of the human bile salt export pump (dissertation). Groningen: Rijks-universiteit Groningen, 2005.
237. Zhao A, Lew JL, Huang L, Yu J, Zhang T, Hrywna Y, *et al.* Human kininogen gene is transactivated by the farnesoid X receptor. *J Biol Chem* 2003 Aug 1;278(31):28765-28770.
238. Qin P, Borges-Marcucci LA, Evans MJ, Harnish DC. Bile acid signaling through FXR induces intracellular adhesion molecule-1 expression in mouse liver and human hepatocytes. *Am J Physiol Gastrointest Liver Physiol* 2005 Aug;289(2):G267-G273.

239. Matsukuma KE, Bennett MK, Huang J, Wang L, Gil G, Osborne TF. Coordinated control of bile acids and lipogenesis through FXR-dependent regulation of fatty acid synthase. *J Lipid Res* 2006 Dec;47(12):2754-2761.
240. Urizar NL, Dowhan DH, Moore DD. The farnesoid X-activated receptor mediates bile acid activation of phospholipid transfer protein gene expression. *J Biol Chem* 2000 Dec 15;275(50):39313-39317.
241. Kast HR, Nguyen CM, Sinal CJ, Jones SA, Laffitte BA, Reue K, *et al.* Farnesoid X-activated receptor induces apolipoprotein C-II transcription: a molecular mechanism linking plasma triglyceride levels to bile acids. *Mol Endocrinol* 2001 Oct;15(10):1720-1728.
242. Huang W, Ma K, Zhang J, Qatanani M, Cuvillier J, Liu J, *et al.* Nuclear receptor-dependent bile acid signaling is required for normal liver regeneration. *Science* 2006 Apr 14;312(5771):233-236.
243. Claudel T, Staels B, Kuipers F. The Farnesoid X receptor: a molecular link between bile acid and lipid and glucose metabolism. *Arterioscler Thromb Vasc Biol* 2005 Oct;25(10):2020-2030.
244. Nguyen A, Bouscarel B. Bile acids and signal transduction: Role in glucose homeostasis. *Cell Signal* 2008 Dec;20(12):2180-97.
245. Id BH, Lagneaux L, Fayyad-Kazan H, Badran B, Najar M, Wiedig M, *et al.* Role of farnesoid X receptor (FXR) in the process of differentiation of bone marrow stromal cells into osteoblasts. *Bone* 2011 Dec;49(6):1219-1231.
246. Huang L, Zhao A, Lew JL, Zhang T, Hrywna Y, Thompson JR, *et al.* Farnesoid X receptor activates transcription of the phospholipid pump MDR3. *J Biol Chem* 2003 Dec 19;278(51):51085-51090.
247. Gnerre C, Blattler S, Kaufmann MR, Looser R, Meyer UA. Regulation of CYP3A4 by the bile acid receptor FXR: evidence for functional binding sites in the CYP3A4 gene. *Pharmacogenetics* 2004 Oct;14(10):635-645.
248. Sanyal S, Bavner A, Haroniti A, Nilsson LM, Lundasen T, Rehnmark S, *et al.* Involvement of corepressor complex subunit GPS2 in transcriptional pathways governing human bile acid biosynthesis. *Proc Natl Acad Sci U S A* 2007 Oct 2;104(40):15665-15670.
249. Lu Y, Heydel JM, Li X, Bratton S, Lindblom T, Radominska-Pandya A. Lithocholic acid decreases expression of UGT2B7 in Caco-2 cells: a potential role for a negative farnesoid X receptor response element. *Drug Metab Dispos* 2005 Jul;33(7):937-946.
250. Kardassis D, Roussou A, Papakosta P, Boulias K, Talianidis I, Zannis VI. Synergism between nuclear receptors bound to specific hormone response elements of the hepatic control region-1 and the proximal apolipoprotein C-II promoter mediate apolipoprotein C-II gene regulation by bile acids and retinoids. *Biochem J* 2003 Jun 1;372(Pt 2):291-304.
251. Hubbert ML, Zhang Y, Lee FY, Edwards PA. Regulation of hepatic Insig-2 by the farnesoid X receptor. *Mol Endocrinol* 2007 Jun;21(6):1359-1369.
252. Chao F, Gong W, Zheng Y, Li Y, Huang G, Gao M, *et al.* Upregulation of scavenger receptor class B type I expression by activation of FXR in hepatocyte. *Atherosclerosis* 2010 Dec;213(2):443-8.
253. Peyer AK, Jung D, Beer M, Gnerre C, Keogh A, Stroka D, *et al.* Regulation of human liver delta-aminolevulinic acid synthase by bile acids. *Hepatology* 2007 Dec;46(6):1960-1970.
254. Lee FY, Kast-Woelbern HR, Chang J, Luo G, Jones SA, Fishbein MC, *et al.* Alpha-crystallin is a target gene of the farnesoid X-activated receptor in human livers. *J Biol Chem* 2005 Sep 9;280(36):31792-31800.
255. Bungard CI, McGivan JD. Identification of the promoter elements involved in the stimulation of ASCT2 expression by glutamine availability in HepG2 cells and the probable involvement of FXR/RXR dimers. *Arch Biochem Biophys* 2005 Nov 15;443(1-2):53-59.
256. Zhang Q, He F, Kuruba R, Gao X, Wilson A, Li J, *et al.* FXR-mediated regulation of angiotensin type 2 receptor expression in vascular smooth muscle cells. *Cardiovasc Res* 2008 Feb 1;77(3):560-569.
257. Hu T, Chouinard M, Cox AL, Sipes P, Marcelo M, Ficorilli J, *et al.* Farnesoid X receptor agonist reduces serum asymmetric dimethylarginine levels through hepatic dimethylarginine dimethylaminohydrolase-1 gene regulation. *J Biol Chem* 2006 Dec 29;281(52):39831-39838.
258. Li J, Wilson A, Kuruba R, Zhang Q, Gao X, He F, *et al.* FXR-mediated regulation of eNOS expression in vascular endothelial cells. *Cardiovasc Res* 2008 Jan;77(1):169-177.

259. Murakami T, Walczak R, Caron S, Duhem C, Vidal V, Darteil R, *et al.* The farnesoid X receptor induces fetuin-B gene expression in human hepatocytes. *Biochem J* 2007 Nov 1;407(3):461-469.
260. Renga B, Mencarelli A, Cipriani S, D'Amore C, Zampella A, Monti MC, *et al.* The nuclear receptor FXR regulates hepatic transport and metabolism of glutamine and glutamate. *Biochim Biophys Acta* 2011 Nov;1812(11):1522-31.
261. Lee H, Hubbert ML, Osborne TF, Woodford K, Zerangue N, Edwards PA. Regulation of the sodium/sulfate co-transporter by farnesoid X receptor alpha. *J Biol Chem* 2007 Jul 27;282(30):21653-21661.
262. Popowski K, Eloranta JJ, Saborowski M, Fried M, Meier PJ, Kullak-Ublick GA. The human organic anion transporter 2 gene is transactivated by hepatocyte nuclear factor-4 alpha and suppressed by bile acids. *Mol Pharmacol* 2005 May;67(5):1629-1638.
263. Jung D, Mangelsdorf DJ, Meyer UA. Pregnane X receptor is a target of farnesoid X receptor. *J Biol Chem* 2006 Jul 14;281(28):19081-19091.
264. Anisfeld AM, Kast-Woelbern HR, Meyer ME, Jones SA, Zhang Y, Williams KJ, *et al.* Syndecan-1 expression is regulated in an isoform-specific manner by the farnesoid-X receptor. *J Biol Chem* 2003 May 30;278(22):20420-20428.
265. Chignard N, Mergely M, Barbu V, Finzi L, Tiret E, Paul A, *et al.* VPAC1 expression is regulated by FXR agonists in the human gallbladder epithelium. *Hepatology* 2005 Sep;42(3):549-557.
266. Kliewer SA, Moore JT, Wade L, Staudinger JL, Watson MA, Jones SA, *et al.* An orphan nuclear receptor activated by pregnanes defines a novel steroid signaling pathway. *Cell* 1998 Jan 9;92(1):73-82.
267. Blumberg B, Sabbagh W, Jr., Juguilon H, Bolado J, Jr., van Meter CM, Ong ES, *et al.* SXR, a novel steroid and xenobiotic-sensing nuclear receptor. *Genes Dev* 1998 Oct 15;12(20):3195-3205.
268. Bertilsson G, Heidrich J, Svensson K, Asman M, Jendeberg L, Sydow-Backman M, *et al.* Identification of a human nuclear receptor defines a new signaling pathway for CYP3A induction. *Proc Natl Acad Sci U S A* 1998 Oct 13;95(21):12208-12213.
269. Staudinger JL, Goodwin B, Jones SA, Hawkins-Brown D, MacKenzie KI, LaTour A, *et al.* The nuclear receptor PXR is a lithocholic acid sensor that protects against liver toxicity. *Proc Natl Acad Sci U S A* 2001 Mar 13;98(6):3369-3374.
270. Xie W, Radominska-Pandya A, Shi Y, Simon CM, Nelson MC, Ong ES, *et al.* An essential role for nuclear receptors SXR/PXR in detoxification of cholestatic bile acids. *Proc Natl Acad Sci U S A* 2001 Mar 13;98(6):3375-3380.
271. Lehmann JM, McKee DD, Watson MA, Willson TM, Moore JT, Kliewer SA. The human orphan nuclear receptor PXR is activated by compounds that regulate CYP3A4 gene expression and cause drug interactions. *J Clin Invest* 1998 Sep 1;102(5):1016-1023.
272. Frank C, Makkonen H, Dunlop TW, Matilainen M, Vaisanen S, Carlberg C. Identification of pregnane X receptor binding sites in the regulatory regions of genes involved in bile acid homeostasis. *J Mol Biol* 2005 Feb 18;346(2):505-519.
273. Synold TW, Dussault I, Forman BM. The orphan nuclear receptor SXR coordinately regulates drug metabolism and efflux. *Nat Med* 2001 May;7(5):584-590.
274. Sonoda J, Xie W, Rosenfeld JM, Barwick JL, Guzelian PS, Evans RM. Regulation of a xenobiotic sulfonation cascade by nuclear pregnane X receptor (PXR). *Proc Natl Acad Sci U S A* 2002 Oct 15;99(21):13801-13806.
275. Teng S, Jekerle V, Piquette-Miller M. Induction of ABCC3 (MRP3) by pregnane X receptor activators. *Drug Metab Dispos* 2003 Nov;31(11):1296-1299.
276. Geick A, Eichelbaum M, Burk O. Nuclear receptor response elements mediate induction of intestinal MDR1 by rifampin. *J Biol Chem* 2001 May 4;276(18):14581-14587.
277. Wistuba W, Gnewuch C, Liebisch G, Schmitz G, Langmann T. Lithocholic acid induction of the FGF19 promoter in intestinal cells is mediated by PXR. *World J Gastroenterol* 2007 Aug 21;13(31):4230-4235.
278. Song KH, Li T, Owsley E, Strom S, Chiang JY. Bile acids activate fibroblast growth factor 19 signaling in human hepatocytes to inhibit cholesterol 7 α -hydroxylase gene expression. *Hepatology* 2009 Jan;49(1):297-305.

279. Inagaki T, Choi M, Moschetta A, Peng L, Cummins CL, McDonald JG, *et al.* Fibroblast growth factor 15 functions as an enterohepatic signal to regulate bile acid homeostasis. *Cell Metab* 2005 Oct;2(4):217-225.
280. Makishima M, Lu T-T, Xie W, Whitfield GK, Domoto H, Evans RM, *et al.* Vitamin D receptor as an intestinal bile acid sensor. *Science* 2002 May 17;296(5571):1313-1316.
281. Adachi R, Honma Y, Masuno H, Kawana K, Shimomura I, Yamada S, *et al.* Selective activation of vitamin D receptor by lithocholic acid acetate, a bile acid derivative. *J Lipid Res* 2005 Jan;46(1):46-57.
282. St-Arnaud R. The direct role of vitamin D on bone homeostasis. *Arch Biochem Biophys* 2008 May 15;473(2):225-230.
283. Carlberg C, Bendik I, Wyss A, Meier E, Sturzenbecker LJ, Grippo JF, *et al.* Two nuclear signalling pathways for vitamin D. *Nature* 1993 Feb 18;361(6413):657-660.
284. Carlberg C, Seuter S. A genomic perspective on vitamin D signaling. *Anticancer Res* 2009 Sep;29(9):3485-3493.
285. Drocourt L, Ourlin JC, Pascussi JM, Maurel P, Vilarem MJ. Expression of CYP3A4, CYP2B6, and CYP2C9 is regulated by the vitamin D receptor pathway in primary human hepatocytes. *J Biol Chem* 2002 Jul 12;277(28):25125-25132.
286. Tachibana S, Yoshinari K, Chikada T, Toriyabe T, Nagata K, Yamazoe Y. Involvement of Vitamin D receptor in the intestinal induction of human ABCB1. *Drug Metab Dispos* 2009 Aug;37(8):1604-1610.
287. McCarthy TC, Li X, Sinal CJ. Vitamin D receptor-dependent regulation of colon multidrug resistance-associated protein 3 gene expression by bile acids. *J Biol Chem* 2005 Jun 17;280(24):23232-23242.
288. Echchgadda I, Song CS, Roy AK, Chatterjee B. Dehydroepiandrosterone sulfotransferase is a target for transcriptional induction by the vitamin D receptor. *Mol Pharmacol* 2004 Mar;65(3):720-729.
289. Chatterjee B, Echchgadda I, Song CS. Vitamin D receptor regulation of the steroid/bile acid sulfotransferase SULT2A1. *Methods Enzymol* 2005;400:165-191.
290. Honjo Y, Sasaki S, Kobayashi Y, Misawa H, Nakamura H. 1,25-dihydroxyvitamin D3 and its receptor inhibit the chenodeoxycholic acid-dependent transactivation by farnesoid X receptor. *J Endocrinol* 2006 Mar;188(3):635-643.
291. Baes M, Gulick T, Choi HS, Martinoli MG, Simha D, Moore DD. A new orphan member of the nuclear hormone receptor superfamily that interacts with a subset of retinoic acid response elements. *Mol Cell Biol* 1994 Mar;14(3):1544-1552.
292. Choi HS, Chung M, Tzamelis I, Simha D, Lee YK, Seol W, *et al.* Differential transactivation by two isoforms of the orphan nuclear hormone receptor CAR. *J Biol Chem* 1997 Sep 19;272(38):23565-23571.
293. Tzamelis I, Moore DD. Role reversal: new insights from new ligands for the xenobiotic receptor CAR. *Trends Endocrinol Metab* 2001 Jan;12(1):7-10.
294. Forman BM, Tzamelis I, Choi HS, Chen J, Simha D, Seol W, *et al.* Androstane metabolites bind to and deactivate the nuclear receptor CAR-beta. *Nature* 1998 Oct 8;395(6702):612-615.
295. Huang W, Zhang J, Chua SS, Qatanani M, Han Y, Granata R, *et al.* Induction of bilirubin clearance by the constitutive androstane receptor (CAR). *Proc Natl Acad Sci U S A* 2003 Apr 1;100(7):4156-4161.
296. Zhang J, Huang W, Qatanani M, Evans RM, Moore DD. The constitutive androstane receptor and pregnane X receptor function coordinately to prevent bile acid-induced hepatotoxicity. *J Biol Chem* 2004 Nov 19;279(47):49517-49522.
297. Guo GL, Lambert G, Negishi M, Ward JM, Brewer HB, Jr., Kliewer SA, *et al.* Complementary roles of farnesoid X receptor, pregnane X receptor, and constitutive androstane receptor in protection against bile acid toxicity. *J Biol Chem* 2003 Nov 14;278(46):45062-45071.
298. Beilke LD, Aleksunes LM, Holland RD, Besselsen DG, Beger RD, Klaassen CD, *et al.* Constitutive androstane receptor-mediated changes in bile acid composition contributes to hepatoprotection from lithocholic acid-induced liver injury in mice. *Drug Metab Dispos* 2009 May;37(5):1035-1045.
299. Saini SP, Sonoda J, Xu L, Toma D, Uppal H, Mu Y, *et al.* A novel constitutive androstane receptor-mediated and CYP3A-independent pathway of bile acid detoxification. *Mol Pharmacol* 2004 Feb;65(2):292-300.

300. Goodwin B, Hodgson E, D'Costa DJ, Robertson GR, Liddle C. Transcriptional regulation of the human CYP3A4 gene by the constitutive androstane receptor. *Mol Pharmacol* 2002 Aug;62(2):359-365.
301. Nitta M, Ku S, Brown C, Okamoto AY, Shan B. CPF: an orphan nuclear receptor that regulates liver-specific expression of the human cholesterol 7 α -hydroxylase gene. *Proc Natl Acad Sci U S A* 1999 Jun 8;96(12):6660-6665.
302. Lu TT, Repa JJ, Mangelsdorf DJ. Orphan nuclear receptors as eLiXIRs and FiXeRs of sterol metabolism. *J Biol Chem* 2001 Oct 12;276(41):37735-37738.
303. Wang ZN, Bassett M, Rainey WE. Liver receptor homologue-1 is expressed in the adrenal and can regulate transcription of 11 beta-hydroxylase. *J Mol Endocrinol* 2001 Oct;27(2):255-258.
304. Ueda H, Sun GC, Murata T, Hirose S. A novel DNA-binding motif abuts the zinc finger domain of insect nuclear hormone receptor FTZ-F1 and mouse embryonal long terminal repeat-binding protein. *Mol Cell Biol* 1992 Dec;12(12):5667-5672.
305. Forman BM. Are those phospholipids in your pocket? *Cell Metab* 2005 Mar;1(3):153-155.
306. Sablin EP, Krylova IN, Fletterick RJ, Ingraham HA. Structural basis for ligand-independent activation of the orphan nuclear receptor LRH-1. *Mol Cell* 2003 Jun;11(6):1575-1585.
307. Lee YK, Parker KL, Choi HS, Moore DD. Activation of the promoter of the orphan receptor SHP by orphan receptors that bind DNA as monomers. *J Biol Chem* 1999 Jul 23;274(30):20869-20873.
308. Del Castillo-Olivares A, Gil G. Alpha 1-fetoprotein transcription factor is required for the expression of sterol 12 α -hydroxylase, the specific enzyme for cholic acid synthesis. Potential role in the bile acid-mediated regulation of gene transcription. *J Biol Chem* 2000 Jun 9;275(23):17793-17799.
309. Chen F, Ma L, Dawson PA, Sinal CJ, Sehayek E, Gonzalez FJ, *et al.* Liver receptor homologue-1 mediates species- and cell line-specific bile acid-dependent negative feedback regulation of the apical sodium-dependent bile acid transporter. *J Biol Chem* 2003 May 30;278(22):19909-19916.
310. Inokuchi A, Hinoshita E, Iwamoto Y, Kohno K, Kuwano M, Uchiumi T. Enhanced expression of the human multidrug resistance protein 3 by bile salt in human enterocytes. A transcriptional control of a plausible bile acid transporter. *J Biol Chem* 2001 Dec 14;276(50):46822-46829.
311. Song X, Kaimal R, Yan B, Deng R. Liver receptor homolog 1 transcriptionally regulates human bile salt export pump expression. *J Lipid Res* 2008 May;49(5):973-984.
312. Frankenberg T, Rao A, Chen F, Haywood J, Shneider BL, Dawson PA. Regulation of the mouse organic solute transporter alpha-beta, Osta-Ost β , by bile acids. *Am J Physiol Gastrointest Liver Physiol* 2006 May;290(5):G912-G922.
313. Freeman LA, Kennedy A, Wu J, Bark S, Remaley AT, Santamarina-Fojo S, *et al.* The orphan nuclear receptor LRH-1 activates the ABCG5/ABCG8 intergenic promoter. *J Lipid Res* 2004 Jul;45(7):1197-1206.
314. Del Castillo-Olivares A, Gil G. Role of FXR and FTF in bile acid-mediated suppression of cholesterol 7 α -hydroxylase transcription. *Nucleic Acids Res* 2000 Sep 15;28(18):3587-3593.
315. Matsukuma KE, Wang L, Bennett MK, Osborne TF. A key role for orphan nuclear receptor liver receptor homologue-1 in activation of fatty acid synthase promoter by liver X receptor. *J Biol Chem* 2007 Jul 13;282(28):20164-20171.
316. Lee YK, Schmidt DR, Cummins CL, Choi M, Peng L, Zhang Y, *et al.* Liver receptor homolog-1 regulates bile acid homeostasis but is not essential for feedback regulation of bile acid synthesis. *Mol Endocrinol* 2008 Jun;22(6):1345-1356.
317. Matakı C, Magnier BC, Houten SM, Annicotte JS, Argmann C, Thomas C, *et al.* Compromised intestinal lipid absorption in mice with a liver-specific deficiency of liver receptor homolog 1. *Mol Cell Biol* 2007 Dec;27(23):8330-8339.
318. Bohan A, Chen WS, Denson LA, Held MA, Boyer JL. Tumor necrosis factor alpha-dependent up-regulation of Lrh-1 and Mrp3(Abcc3) reduces liver injury in obstructive cholestasis. *J Biol Chem* 2003 Sep 19;278(38):36688-36698.
319. Chong HK, Biesinger J, Seo YK, Xie X, Osborne TF. Genome-wide analysis of hepatic LRH-1 reveals a promoter binding preference and suggests a role in regulating genes of lipid metabolism in concert with FXR. *BMC Genomics* 2012;13:51.

320. Yang Y, Zhang M, Eggertsen G, Chiang JY. On the mechanism of bile acid inhibition of rat sterol 12 α -hydroxylase gene (CYP8B1) transcription: roles of alpha-fetoprotein transcription factor and hepatocyte nuclear factor 4 α . *Biochim Biophys Acta* 2002 Jun 13;1583(1):63-73.
321. Miao J, Choi SE, Seok SM, Yang L, Zuercher WJ, Xu Y, *et al*. Ligand-dependent regulation of the activity of the orphan nuclear receptor, small heterodimer partner (SHP), in the repression of bile acid biosynthetic CYP7A1 and CYP8B1 genes. *Mol Endocrinol* 2011 Jul;25(7):1159-1169.
322. Lee HK, Lee YK, Park SH, Kim YS, Park SH, Lee JW, *et al*. Structure and expression of the orphan nuclear receptor SHP gene. *J Biol Chem* 1998 Jun 5;273(23):14398-14402.
323. Oropeza CE, Li L, McLachlan A. Differential inhibition of nuclear hormone receptor dependent hepatitis B virus replication by small heterodimer partner. *J Virol* 2008 Apr;82(8):3814-21.
324. He N, Park K, Zhang Y, Huang J, Lu S, Wang L. Epigenetic inhibition of nuclear receptor small heterodimer partner is associated with and regulates hepatocellular carcinoma growth. *Gastroenterology* 2008 Mar;134(3):793-802.
325. Volle DH, Duggavathi R, Magnier BC, Houten SM, Cummins CL, Lobaccaro JM, *et al*. The small heterodimer partner is a gonadal gatekeeper of sexual maturation in male mice. *Genes Dev* 2007 Feb 1;21(3):303-315.
326. Nishigori H, Tomura H, Tonooka N, Kanamori M, Yamada S, Shio K, *et al*. Mutations in the small heterodimer partner gene are associated with mild obesity in Japanese subjects. *Proc Natl Acad Sci U S A* 2001 Jan 16;98(2):575-580.
327. Mitchell SM, Weedon MN, Owen KR, Shields B, Wilkins-Wall B, Walker M, *et al*. Genetic variation in the small heterodimer partner gene and young-onset type 2 diabetes, obesity, and birth weight in U.K. subjects. *Diabetes* 2003 May;52(5):1276-1279.
328. Goodwin B, Watson MA, Kim H, Miao J, Kemper JK, Kliewer SA. Differential regulation of rat and human CYP7A1 by the nuclear oxysterol receptor liver X receptor- α . *Mol Endocrinol* 2003 Mar;17(3):386-394.
329. Seol W, Choi HS, Moore DD. An orphan nuclear hormone receptor that lacks a DNA binding domain and heterodimerizes with other receptors. *Science* 1996 May 31;272(5266):1336-1339.
330. Willy PJ, Umesono K, Ong ES, Evans RM, Heyman RA, Mangelsdorf DJ. LXR, a nuclear receptor that defines a distinct retinoid response pathway. *Genes Dev* 1995 May 1;9(9):1033-1045.
331. Apfel R, Benbrook D, Lernhardt E, Ortiz MA, Salbert G, Pfahl M. A novel orphan receptor specific for a subset of thyroid hormone-responsive elements and its interaction with the retinoid/thyroid hormone receptor subfamily. *Mol Cell Biol* 1994 Oct;14(10):7025-7035.
332. Song C, Kokontis JM, Hiipakka RA, Liao S. Ubiquitous receptor: a receptor that modulates gene activation by retinoic acid and thyroid hormone receptors. *Proc Natl Acad Sci U S A* 1994 Nov 8;91(23):10809-10813.
333. Forman BM, Ruan B, Chen J, Schroepfer GJ, Jr., Evans RM. The orphan nuclear receptor LXR α is positively and negatively regulated by distinct products of mevalonate metabolism. *Proc Natl Acad Sci U S A* 1997 Sep 30;94(20):10588-10593.
334. Lehmann JM, Kliewer SA, Moore LB, Smith-Oliver TA, Oliver BB, Su JL, *et al*. Activation of the nuclear receptor LXR by oxysterols defines a new hormone response pathway. *J Biol Chem* 1997 Feb 7;272(6):3137-3140.
335. Song C, Hiipakka RA, Liao S. Selective activation of liver X receptor α by 6 α -hydroxy bile acids and analogs. *Steroids* 2000 Aug;65(8):423-427.
336. Janowski BA, Willy PJ, Devi TR, Falck JR, Mangelsdorf DJ. An oxysterol signalling pathway mediated by the nuclear receptor LXR α . *Nature* 1996 Oct 24;383(6602):728-731.
337. Dahlman-Wright K, Zhao C. LXR in cholesterol metabolism. *J Endocrinol* 2010 Mar;204(3):233-40.
338. Ulven SM, Dalen KT, Gustafsson JA, Nebb HI. LXR is crucial in lipid metabolism. *Prostaglandins Leukot Essent Fatty Acids* 2005 Jul;73(1):59-63.
339. Peet DJ, Turley SD, Ma W, Janowski BA, Lobaccaro JM, Hammer RE, *et al*. Cholesterol and bile acid metabolism are impaired in mice lacking the nuclear oxysterol receptor LXR α . *Cell* 1998 May 29;93(5):693-704.

340. Edwards PA, Kast HR, Anisfeld AM. BAREing it all: the adoption of LXR and FXR and their roles in lipid homeostasis. *J Lipid Res* 2002 Jan;43(1):2-12.
341. Repa JJ, Berge KE, Pomajzl C, Richardson JA, Hobbs H, Mangelsdorf DJ. Regulation of ATP-binding cassette sterol transporters ABCG5 and ABCG8 by the liver X receptors alpha and beta. *J Biol Chem* 2002 May 24;277(21):18793-18800.
342. Pare JF, Malenfant D, Courtemanche C, Jacob-Wagner M, Roy S, Allard D, *et al.* The fetoprotein transcription factor (FTF) gene is essential to embryogenesis and cholesterol homeostasis and is regulated by a DR4 element. *J Biol Chem* 2004 May 14;279(20):21206-21216.
343. Landrier JF, Grober J, Demydchuk J, Besnard P. FXRE can function as an LXRE in the promoter of human ileal bile acid-binding protein (I-BABP) gene. *FEBS Lett* 2003 Oct 23;553(3):299-303.
344. Okuwaki M, Takada T, Iwayanagi Y, Koh S, Kariya Y, Fujii H, *et al.* LXR alpha transactivates mouse organic solute transporter alpha and beta via IR-1 elements shared with FXR. *Pharm Res* 2007 Feb;24(2):390-398.
345. Wang J, Einarsson C, Murphy C, Parini P, Bjorkhem I, Gafvels M, *et al.* Studies on LXR- and FXR-mediated effects on cholesterol homeostasis in normal and cholic acid-depleted mice. *J Lipid Res* 2006 Feb;47(2):421-430.
346. Wang L, Lee YK, Bundman D, Han Y, Thevananther S, Kim CS, *et al.* Redundant pathways for negative feedback regulation of bile acid production. *Dev Cell* 2002 Jun;2(6):721-731.
347. Desvergne B. RXR: from partnership to leadership in metabolic regulations. *Vitam Horm* 2007;75:1-32.
348. Issemann I, Prince RA, Tugwood JD, Green S. The retinoid X receptor enhances the function of the peroxisome proliferator activated receptor. *Biochimie* 1993;75(3-4):251-256.
349. Kliewer SA, Umesono K, Noonan DJ, Heyman RA, Evans RM. Convergence of 9-cis retinoic acid and peroxisome proliferator signalling pathways through heterodimer formation of their receptors. *Nature* 1992 Aug 27;358(6389):771-774.
350. DiRenzo J, Soderstrom M, Kurokawa R, Ogliastro MH, Ricote M, Ingrey S, *et al.* Peroxisome proliferator-activated receptors and retinoic acid receptors differentially control the interactions of retinoid X receptor heterodimers with ligands, coactivators, and corepressors. *Mol Cell Biol* 1997 Apr;17(4):2166-2176.
351. Forman BM, Umesono K, Chen J, Evans RM. Unique response pathways are established by allosteric interactions among nuclear hormone receptors. *Cell* 1995 May 19;81(4):541-550.
352. Ferrara J, McCuaig K, Hendy GN, Uskokovic M, White JH. Highly potent transcriptional activation by 16-ene derivatives of 1,25-dihydroxyvitamin D₃. Lack of modulation by 9-cis-retinoic acid of response to 1,25-dihydroxyvitamin D₃ or its derivatives. *J Biol Chem* 1994 Jan 28;269(4):2971-2981.
353. MacDonald PN, Dowd DR, Nakajima S, Galligan MA, Reeder MC, Haussler CA, *et al.* Retinoid X receptors stimulate and 9-cis retinoic acid inhibits 1,25-dihydroxyvitamin D₃-activated expression of the rat osteocalcin gene. *Mol Cell Biol* 1993 Sep;13(9):5907-5917.
354. Castillo AI, Sanchez-Martinez R, Moreno JL, Martinez-Iglesias OA, Palacios D, Aranda A. A permissive retinoid X receptor/thyroid hormone receptor heterodimer allows stimulation of prolactin gene transcription by thyroid hormone and 9-cis-retinoic acid. *Mol Cell Biol* 2004 Jan;24(2):502-513.
355. Tzamelis I, Chua SS, Cheskis B, Moore DD. Complex effects of rexinoids on ligand dependent activation or inhibition of the xenobiotic receptor, CAR. *Nucl Recept* 2003 Jun 6;1(1):2.
356. Kassam A, Miao B, Young PR, Mukherjee R. Retinoid X receptor (RXR) agonist-induced antagonism of farnesoid X receptor (FXR) activity due to absence of coactivator recruitment and decreased DNA binding. *J Biol Chem* 2003 Mar 21;278(12):10028-10032.
357. Del Castillo-Olivares A, Gil G. Suppression of sterol 12 α -hydroxylase transcription by the short heterodimer partner: insights into the repression mechanism. *Nucleic Acids Res* 2001 Oct 1;29(19):4035-4042.
358. Li H, Chen F, Shang Q, Pan L, Shneider BL, Chiang JY, *et al.* FXR-activating ligands inhibit rabbit ASBT expression via FXR-SHP-FTF cascade. *Am J Physiol Gastrointest Liver Physiol* 2005 Jan;288(1):G60-G66.
359. Lee YK, Moore DD. Dual mechanisms for repression of the monomeric orphan receptor liver receptor homologous protein-1 by the orphan small heterodimer partner. *J Biol Chem* 2002 Jan 25;277(4):2463-2467.

- 360. Seol W, Chung M, Moore DD. Novel receptor interaction and repression domains in the orphan receptor SHP. *Mol Cell Biol* 1997 Dec;17(12):7126-7131.
- 361. Kerr TA, Saeki S, Schneider M, Schaefer K, Berdy S, Redder T, *et al.* Loss of nuclear receptor SHP impairs but does not eliminate negative feedback regulation of bile acid synthesis. *Dev Cell* 2002 Jun;2(6):713-720.
- 362. Wang L, Han Y, Kim CS, Lee YK, Moore DD. Resistance of SHP-null mice to bile acid-induced liver damage. *J Biol Chem* 2003 Nov 7;278(45):44475-44481.
- 363. Park YJ, Qatanani M, Chua SS, LaRey JL, Johnson SA, Watanabe M, *et al.* Loss of orphan receptor small heterodimer partner sensitizes mice to liver injury from obstructive cholestasis. *Hepatology* 2008 May;47(5):1578-1586.
- 364. Chiang JY, Kimmel R, Stroup D. Regulation of cholesterol 7 α -hydroxylase gene (CYP7A1) transcription by the liver orphan receptor (LXR α). *Gene* 2001 Jan 10;262(1-2):257-265.
- 365. Agellon LB, Drover VA, Cheema SK, Gbaguidi GF, Walsh A. Dietary cholesterol fails to stimulate the human cholesterol 7 α -hydroxylase gene (CYP7A1) in transgenic mice. *J Biol Chem* 2002 Jun 7;277(23):20131-20134.
- 366. Karpen SJ, Sun AQ, Kudish B, Hagenbuch B, Meier PJ, Ananthanarayanan M, *et al.* Multiple factors regulate the rat liver basolateral sodium-dependent bile acid cotransporter gene promoter. *J Biol Chem* 1996 Jun 21;271(25):15211-15221.
- 367. Trauner M, Arrese M, Lee H, Boyer JL, Karpen SJ. Endotoxin downregulates rat hepatic ntcp gene expression via decreased activity of critical transcription factors. *J Clin Invest* 1998 May 15;101(10):2092-2100.
- 368. Jung D, Hagenbuch B, Fried M, Meier PJ, Kullak-Ublick GA. Role of liver-enriched transcription factors and nuclear receptors in regulating the human, mouse, and rat NTCP gene. *Am J Physiol Gastrointest Liver Physiol* 2004 May;286(5):G752-G761.
- 369. Jung D, Kullak-Ublick GA. Hepatocyte nuclear factor 1 α : a key mediator of the effect of bile acids on gene expression. *Hepatology* 2003 Mar;37(3):622-631.
- 370. Eloranta JJ, Jung D, Kullak-Ublick GA. The human Na⁺-taurocholate cotransporting polypeptide gene is activated by glucocorticoid receptor and peroxisome proliferator-activated receptor- γ coactivator-1 α , and suppressed by bile acids via a small heterodimer partner-dependent mechanism. *Mol Endocrinol* 2006 Jan;20(1):65-79.
- 371. Jung D, Fantin AC, Scheurer U, Fried M, Kullak-Ublick GA. Human ileal bile acid transporter gene ASBT (SLC10A2) is transactivated by the glucocorticoid receptor. *Gut* 2004 Jan;53(1):78-84.
- 372. Del Castillo-Olivares A, Campos JA, Pandak WM, Gil G. The role of α -fetoprotein transcription factor/LRH-1 in bile acid biosynthesis: a known nuclear receptor activator that can act as a suppressor of bile acid biosynthesis. *J Biol Chem* 2004 Apr 16;279(16):16813-16821.
- 373. Gupta S, Stravitz RT, Dent P, Hylemon PB. Down-regulation of cholesterol 7 α -hydroxylase (CYP7A1) gene expression by bile acids in primary rat hepatocytes is mediated by the c-Jun N-terminal kinase pathway. *J Biol Chem* 2001 May 11;276(19):15816-15822.
- 374. Jung D, Inagaki T, Gerard RD, Dawson PA, Kliewer SA, Mangelsdorf DJ, *et al.* FXR agonists and FGF15 reduce fecal bile acid excretion in a mouse model of bile acid malabsorption. *J Lipid Res* 2007 Dec;48(12):2693-2700.
- 375. Sinha J, Chen F, Miloh T, Burns RC, Yu Z, Shneider BL. β -Klotho and FGF-15/19 inhibit the apical sodium-dependent bile acid transporter in enterocytes and cholangiocytes. *Am J Physiol Gastrointest Liver Physiol* 2008 Sep 4.
- 376. Jiang W, Miyamoto T, Kakizawa T, Nishio SI, Oiwa A, Takeda T, *et al.* Inhibition of LXR α signaling by vitamin D receptor: possible role of VDR in bile acid synthesis. *Biochem Biophys Res Commun* 2006 Dec 8;351(1):176-184.
- 377. Chiang JY. Regulation of bile acid synthesis: pathways, nuclear receptors, and mechanisms. *J Hepatol* 2004 Mar;40(3):539-551.
- 378. Davis RA, Miyake JH, Hui TY, Spann NJ. Regulation of cholesterol-7 α -hydroxylase: BAREly missing a SHP. *J Lipid Res* 2002 Apr;43(4):533-543.

- 379. Miyake JH, Wang SL, Davis RA. Bile acid induction of cytokine expression by macrophages correlates with repression of hepatic cholesterol 7 α -hydroxylase. *J Biol Chem* 2000 Jul 21;275(29):21805-21808.
- 380. Triantis V, Saeland E, Bijl N, Oude-Elferink RP, Jansen PL. Glycosylation of fibroblast growth factor receptor 4 is a key regulator of fibroblast growth factor 19-mediated down-regulation of cytochrome P450 7A1. *Hepatology* 2010 Aug;52(2):656-666.
- 381. Tomiyama K, Maeda R, Urakawa I, Yamazaki Y, Tanaka T, Ito S, *et al.* Relevant use of Klotho in FGF19 subfamily signaling system *in vivo*. *Proc Natl Acad Sci U S A* 2010 Jan 26;107(4):1666-1671.
- 382. Stravitz RT, Rao YP, Vlahcevic ZR, Gurley EC, Jarvis WD, Hylemon PB. Hepatocellular protein kinase C activation by bile acids: implications for regulation of cholesterol 7 α -hydroxylase. *Am J Physiol* 1996 Aug;271(2 Pt 1):G293-G303.
- 383. Li T, Jahan A, Chiang JY. Bile acids and cytokines inhibit the human cholesterol 7 α -hydroxylase gene via the JNK/c-jun pathway in human liver cells. *Hepatology* 2006 Jun;43(6):1202-1210.
- 384. Gineste R, Sirvent A, Paumelle R, Helleboid S, Aquilina A, Darteil R, *et al.* Phosphorylation of farnesoid X receptor by protein kinase C promotes its transcriptional activity. *Mol Endocrinol* 2008 Nov;22(11):2433-2447.